

On the Sensitivity Complexity of k -Uniform Hypergraph Properties

Qian Li¹ and Xiaoming Sun¹

¹Institute of Computing Technology, Chinese Academy of Sciences,
Beijing, China
{liqian, sunxiaoming}@ict.ac.cn

Abstract

In this paper we investigate the sensitivity complexity of hypergraph properties. We present a k -uniform hypergraph property with sensitivity complexity $O(n^{\lceil k/3 \rceil})$ for any $k \geq 3$, where n is the number of vertices. Moreover, we can do better when $k \equiv 1 \pmod{3}$ by presenting a k -uniform hypergraph property with sensitivity $O(n^{\lceil k/3 \rceil - 1/2})$. This result disproves a conjecture of Babai [1], which conjectures that the sensitivity complexity of k -uniform hypergraph properties is at least $\Omega(n^{k/2})$. We also investigate the sensitivity complexity of other symmetric functions and show that for many classes of transitive Boolean functions the minimum achievable sensitivity complexity can be $O(N^{1/3})$, where N is the number of variables. Finally, we give a lower bound for sensitivity of k -uniform hypergraph properties, which implies the *sensitivity conjecture* of k -uniform hypergraph properties for any constant k .

1 Introduction

In order to understand the effect of symmetry on computational complexity, especially in the decision tree model, Boolean functions with certain symmetry have been extensively investigated. It is observed that symmetry usually implies high complexity or makes the problem harder in the decision tree model. An illustrative example is the well known *evasiveness conjecture*, which asserts that any monotone transitive Boolean function is evasive, and it has attracted a lot of attention [2–5]. Rivest and Vuillemin [6] showed that any non-constant monotone graph property are weakly evasive. Kulkarni et al. [7] showed the analogous result for 3-hypergraph properties. Black [8] extended these results to k -uniform hypergraph properties for any fixed k .

Sensitivity complexity is an important complexity measure of a Boolean function in the decision tree model, and sensitivity complexity of Boolean functions

with certain symmetry has also attracted a lot of attention. One of the most challenging problem here is whether *symmetry* implies high sensitivity complexity. The famous sensitivity conjecture, which asserts sensitivity complexity and block sensitivity are polynomially related, implies $s(f) = \Omega(n^\alpha)$ for transitive functions with some constant $\alpha > 0$ since it has been shown that $bs(f) = \Omega(n^{1/3})$ for transitive functions [9]. Turan [10] initiated the study of sensitivity of graph properties and proved that the sensitivity is greater than $n/4$ for any nontrivial graph property, where n is the number of vertices, and this relation is also tight up to a constant factor. He also pointed out that for symmetric functions, $s(f) \geq n/2 \geq bs(f)/2$. Recently Sun improved the lower bound to $\frac{6}{17}n$ [11], and Gao et al. [12] investigated the sensitivity of bipartite graph properties as well. In 2005, Chakraborty [13] constructed a minterm cyclically invariant Boolean function whose sensitivity is $\Theta(n^{1/3})$, which answers Turan's question [10] in the negative. He also showed this bound is tight for minterm transitive functions.

For hypergraph properties, Biderman et al. [1] present a sequence of k -uniform hypergraph properties with sensitivity $\Theta(\sqrt{N})$, where $N = \binom{n}{k}$ is the number of variables. Babai conjectures that this bound is tight, i.e., $s(f) = \Omega(\sqrt{N})$ for any nontrivial k -uniform hypergraph property f .

Our Results. In this paper we disprove this conjecture by constructing k -uniform hypergraph properties with sensitivity $O(n^{\lceil k/3 \rceil})$, i.e.,

Theorem 1. *For any $k \geq 3$, there exist a sequence of k -uniform hypergraph properties f such that $s(f) = O(n^{\lceil k/3 \rceil})$.*

Moreover, we can give better constructions when $k \equiv 1 \pmod{3}$.

Theorem 2. *For any $k \geq 4$ satisfying $k \equiv 1 \pmod{3}$, there exist a sequence of k -uniform hypergraph properties f such that $s(f) = O(n^{\lceil k/3 \rceil - 1/2})$.*

More generally, we also investigate the sensitivity of k -partite k -uniform hypergraph properties. Actually, the constructions of k -uniform hypergraph properties are inspired by the constructions of k -partite k -uniform hypergraph properties.

Theorem 3. *For any $k \geq 3$, there exist a sequence of k -partite k -uniform hypergraph properties $f : \{0, 1\}^{n^k} \rightarrow \{0, 1\}$ such that $s(f) = O(n^{\lceil k/3 \rceil})$.*

Theorem 4. *For any $k \geq 4$ satisfying $k \equiv 1 \pmod{3}$, there exist a sequence of k -partite k -uniform hypergraph properties $f : \{0, 1\}^{n^k} \rightarrow \{0, 1\}$ such that $s(f) = O(n^{\lceil k/3 \rceil - 1/2})$.*

Let G be an Abelian group, the fundamental theorem of finite abelian groups states that $G \cong C_{m_1} \times \cdots \times C_{m_l}$, where C_m is the cyclic group of order m and $|G| = \prod_{i=1}^l m_i$.

Theorem 5. *Let $G \leq S_n$ be a transitive Abelian group, then there exists a Boolean function $f : \{0, 1\}^n \rightarrow \{0, 1\}$ invariant under G such that $s(f) \leq an^{1/3}$, where a is a number only depending on l .*

On the other side, we prove a lower bound of the sensitivity of k -uniform hypergraph properties, which implies the sensitivity conjecture of k -uniform hypergraph properties.

Theorem 6. *For any constant k and any non-trivial k -uniform hypergraph property f , $s(f) = \Omega(n)$.*

Similar lower bound holds for the sensitivity of k -partite k -uniform hypergraph properties.

Theorem 7. *For any constant k and any non-trivial k -partite k -uniform hypergraph property f , $s(f) = \Omega(n)$, where n is the number of vertices in one partition.*

The proof of this theorem is very similar with the proof of Theorem 2 in [12], except that we divide k -partitions into two sets of size 1 and $k - 1$ respectively first. We omit the proof in this paper.

Related Work. Sensitivity complexity and block sensitivity are first introduced by Cook, Dwork and Reischuk [14, 15] and Nisan [16] respectively, to study the time complexity of CREW-PRAMs. Block sensitivity has been shown to be polynomially related to a number of other complexity measures [17], such as decision tree complexity, certificate complexity, polynomial degree and quantum query complexity, etc, except sensitivity. The famous sensitivity conjecture, proposed by Nisan and Szegedy [18], asserts that block sensitivity and sensitivity complexity are also polynomially related. On one side, it is easy to see $s(f) \leq bs(f)$ for any Boolean function f according to the definitions. On the other side, it is much more challenging to prove or disprove block sensitivity is polynomially bounded by sensitivity. Despite of a lot of effort, the best known upper bound is exponential: $bs(f) \leq \max\{2^{s(f)-1}(s(f) - \frac{1}{3}), s(f)\}$ [19]. Recently, He, Li and Sun further improve the upper bound to $(\frac{8}{9} + o(1))s(f)2^{s(f)-1}$ [20]. The best known separation between sensitivity and block sensitivity is quadratic [21]: there exist a sequence of Boolean functions f with $bs(f) = \frac{2}{3}s(f)^2 - \frac{1}{3}s(f)$. For an excellent survey on the sensitivity conjecture, see [22]. For other recent progress, see [23–34].

Organization. We present some preliminaries in Section 2, and give the proofs of Theorem 1 and Theorem 2 in Section 3. We give the constructions of k -partite k -uniform hypergraph properties (Theorem 3 and 4) and the proof of Theorem 5 in Section 4 and give the proof of Theorem 6 in Section 5. Finally, we conclude this paper with some open problems in Section 6.

2 Preliminaries

Let $f : \{0, 1\}^n \rightarrow \{0, 1\}$ be a Boolean function and $[n] = \{1, 2, \dots, n\}$. For an input $x \in \{0, 1\}^n$ and a subset $B \subseteq [n]$, x^B denotes the input obtained by flipping all the bit x_j such that $j \in B$.

Definition 1. The sensitivity of f on input x is defined as $s(f, x) := |\{i | f(x) \neq f(x^{[i]})\}|$. The sensitivity, 0-sensitivity and 1-sensitivity of the function f are defined as $s(f) := \max_x s(f, x)$, $s_0(f) = \max_{x \in f^{-1}(0)} s(f, x)$ and $s_1(f) = \max_{x \in f^{-1}(1)} s(f, x)$ respectively.

Definition 2. The block sensitivity $bs(f, x)$ of f on input x is the maximum number of disjoint subsets B_1, B_2, \dots, B_r of $[n]$ such that for all $j \in [r]$, $f(x) \neq f(x^{B_j})$. The block sensitivity of f is defined as $bs(f) = \max_x bs(f, x)$.

Definition 3. A partial assignment is a function $p : [n] \rightarrow \{0, 1, \star\}$. We call $S = \{i | p_i \neq \star\}$ the support of this partial assignment. We define the size of p denoted by $|p|$ to be $|S|$. We call x a (full) assignment if $x : [n] \rightarrow \{0, 1\}$. We say x is consistent with p if $x|_S = p$, i.e., $x_i = p_i$ for all $i \in S$.¹

Definition 4. For $b \in \{0, 1\}$, a b -certificate for f is a partial assignment p such that $f(x) = b$ whenever x is consistent with p .

The certificate complexity $C(f, x)$ of f on input x is the minimum size of $f(x)$ -certificate that is consistent with x . The certificate complexity of f is $C(f) = \max_x C(f, x)$.

The 1-certificate complexity of f is $C_1(f) = \max_{x \in f^{-1}(1)} C(f, x)$, and similarly we define $C_0(f)$.

According to the definitions, it's easy to see $s(f) \leq bs(f) \leq C(f)$, $s_0(f) \leq C_0(f)$ and $s_1(f) \leq C_1(f)$.

Definition 5. Let p and p' be two partial assignments, the distance between p and p' is defined as $dist(p, p') = |\{i | p_i = 1 \text{ and } p'_i = 0, \text{ or } p_i = 0 \text{ and } p'_i = 1\}|$.

Definition 6. Let $f : \{0, 1\}^n \rightarrow \{0, 1\}$ be a Boolean function and G be a subgroup of S_n , we say that f is invariant under G if $f(x_1, \dots, x_n) = f(x_{\sigma(1)}, \dots, x_{\sigma(n)})$ for any $x \in \{0, 1\}^n$ and any $\sigma \in G$.

A Boolean function f is called transitive (or weakly symmetric) if G is a transitive group². A Boolean function f is called symmetric if $G = S_n$.

Definition 7. A transitive Boolean function f is called minterm-transitive if there exist a partial assignment p such that $f(x) = 1$ if and only if x is consistent with $p^\sigma := (p_{\sigma(1)}, p_{\sigma(2)}, \dots, p_{\sigma(n)})$ for some $\sigma \in G$. We call p the minterm.

A Boolean string can represent a graph in the following manner: $x_{(i,j)} = 1$ means there is an edge connecting vertex i and vertex j , and $x_{i,j} = 0$ means there is no such edge. Graph properties are functions which are independent with the labeling of vertices, i.e. two isomorphic graphs have the same function value.

Definition 8. A Boolean function $f : \{0, 1\}^{\binom{n}{2}} \rightarrow \{0, 1\}$ is called a graph property if for every input $x = (x_{(1,2)}, \dots, x_{(n-1,n)})$ and every permutation $\sigma \in S_n$,

$$f(x_{(1,2)}, \dots, x_{(n-1,n)}) = f(x_{(\sigma(1), \sigma(2))}, \dots, x_{(\sigma(n-1), \sigma(n))}).$$

¹ The function p can be viewed as a vector, and we sometimes use p_i to represent $p(i)$.

² A group $G \leq S_n$ is transitive if for every $i < j$, there exists a $\sigma \in G$ such that $\sigma(i) = j$.

Similarly, we define k -uniform hypergraph properties.

Definition 9. A Boolean function $f : \{0, 1\}^{\binom{[n]}{k}} \rightarrow \{0, 1\}$ is called a k -uniform hypergraph property if for every input $x = (x_{(1,2,\dots,k)}, \dots, x_{(n-k+1,\dots,n)})$ and every permutation $\sigma \in S_n$,

$$f(x_{(1,2,\dots,k)}, \dots, x_{(n-k+1,\dots,n)}) = f(x_{(\sigma(1),\sigma(2),\dots,\sigma(k))}, \dots, x_{(\sigma(n-k+1),\dots,\sigma(n))}).$$

Let p be a partial assignment and $\sigma \in S_n$, we define $\sigma(p)$ as $\sigma(p)_S = p_{\sigma(S)}$ where S is any subset of $[n]$ of size k and $\sigma(S) = \{\sigma(i) | i \in S\}$.

Definition 10. A Boolean function $f : \{0, 1\}^{n^k} \rightarrow \{0, 1\}$ is called k -partite k -uniform hypergraph property, if for every input $x = (x_{(1,1,\dots,1)}, \dots, x_{(n,n,\dots,n)})$ and every $\sigma = (\sigma_1, \dots, \sigma_k) \in S_n^{\otimes k}$,

$$f(x_{(1,1,\dots,1)}, \dots, x_{(n,n,\dots,n)}) = f(x_{(\sigma_1(1),\dots,\sigma_k(1))}, \dots, x_{(\sigma_1(n),\dots,\sigma_k(n))}).$$

It is easy to see that any (k -partite) k -uniform hypergraph property is transitive.

3 k -Uniform Hypergraph Properties

In this section, we give the proofs of Theorem 1 and Theorem 2.

Proof. (Proof of Theorem 1) The function we construct is a minterm function. Let p be the minterm defining f , and it is constructed as follow:

First, let k_1 and k_2 be two integers such that $k_1 + 2k_2 = k$ and $k_1, k_2 \leq \lceil k/3 \rceil$. Let $V = \{v_1, \dots, v_n\}$ be the set of vertices and $B = \{v_n, v_{n-1}, \dots, v_{n-k_1+1}\}$. For each $1 \leq i \leq 6$, let $W_i = \{v_{(i-1)k_2+1}, \dots, v_{ik_2}\}$, and $C = \bigcup_{1 \leq i \leq 6} W_i$, $D = V \setminus (C \cup B)$.

- For any $S \subseteq C$ of size $2k_2$, $p(B \cup S) = 0$, except $S = W_i \cup W_{i+1}$ for $i \in [5]$ where $p(B \cup S) = 1$.
- For any S of size $2k_2$ and $k_2 \leq |S \cap C| < 2k_2$, $p(B \cup S) = 1$, except W_3 or $W_4 \subseteq S$ where $p(B \cup S) = 0$.
- All the other variables are \star .

If $f(x) = 1$ then x is consistent with some $\sigma(p)$, which implies $C(f, x) \leq |p|$. Thus $s_1(f) \leq C_1(f) \leq |p| = \sum_{i=k_2}^{2k_2} \binom{6k_2}{i} \binom{n-6k_2-k_1}{2k_2-i} = O(n^{k_2})$. Moreover, if $f(x) = 0$ then $s(f, x)$ is at most the number of isomorphisms of p (i.e., $\sigma(p)$ s) adjacent to x , thus according to the triangle inequality, $s_0(f)$ is at most the maximum number of $\sigma(p)$ s where the distance between any two of them is at most 2. We claim that for any $\pi(p)$, there are $O(1)$ isomorphisms $\sigma(p)$ s satisfying $\pi(B) = \sigma(B)$ and $\text{dist}(\pi(p), \sigma(p)) \leq 2$. It is easy to see that this claim implies $s_0(f) = O(n^{k_1})$ since there are $\binom{n}{k_1} = O(n^{k_1})$ possible choices of the $\sigma(B)$ s, and this will end the whole proof.




Figure 1: The graph to illustrate p for $k_2 = 1$

Claim 1. For any $\pi(p)$, there are only $O(1)$ $\sigma(p)$ s satisfying $\pi(B) = \sigma(B)$ and $\text{dist}(\pi(p), \sigma(p)) \leq 2$.

Proof. It is easy to see that this claim is equivalent to show $|\{\sigma(p) | \text{dist}(p, \sigma(p)) \leq 2 \text{ and } \sigma(B) = B\}| = O(1)$. The case for $k_2 = 1$ is a little special, and we discuss this case first.

Case for $k_2 = 1$ We use Figure 1 to illustrate p . Note that the vertices in D are symmetric and $|C| = O(1)$, thus $|\{\sigma(p) | \sigma(C) = C \text{ and } \sigma(B) = B\}| = O(1)$. So we only need to consider the set $\{\sigma(p) | \sigma(C) \neq C \text{ and } \sigma(B) = B\}$, and we exclude each σ case by case:

1. $\sigma(W_3)$ or $\sigma(W_4) \in \{W_1, W_2, W_5, W_6\}$.

W.L.O.G, assume $\sigma(W_3) = W_1$, then

$$\begin{aligned}
 \text{dist}(p, \sigma(p)) &\geq |\{e \subseteq [n] | \sigma(p)(e) = 1, p(e) = 0, |e| = k, \{W_3, B\} \subseteq e\}| \\
 &\geq |\{e \subseteq [n] | \sigma(p)(e) = 1, |e| = k, \{W_3, B\} \subseteq e\}| \\
 &\quad - |\{e \subseteq [n] | p(e) = \{1, \star\}, |e| = k, \{W_3, B\} \subseteq e\}| \\
 &= |\{e \subseteq [n] | p(e) = 1, |e| = k, \{W_1, B\} \subseteq e\}| - O(1) \\
 &\geq n - O(1) \geq 3.
 \end{aligned}$$

2. $\sigma(W_3)$ or $\sigma(W_4) \in D$, and $\{\sigma(W_3), \sigma(W_4)\} \cap \{W_1, W_2, W_5, W_6\} = \emptyset$.

W.L.O.G, assume $\sigma(W_3) \in D$, note that for any v , $p(B \cup W_3 \cup v) \neq \star$, and $|\{v \neq W_4 | p(B \cup W_3 \cup v) = 1\}| = 1$. While $|\{v \neq W_4 | \sigma(p)(B \cup W_3 \cup v) = 1\}| = |\{v \neq W_4 | p(B \cup \sigma(W_3) \cup \sigma(v)) = 1\}| = 4$, thus $\text{dist}(p, \sigma(p)) \geq 3$.

3. $\sigma(W_3) = W_3$ and $\sigma(W_4) = W_4$.

- (a) $\sigma(W_5) \neq W_5$ and $\sigma(W_2) \neq W_2$.

Since $p(B \cup \sigma(W_2) \cup \sigma(W_3)) = p(B \cup \sigma(W_4) \cup \sigma(W_5)) = 0$ and $p(B \cup \sigma(W_3) \cup \sigma(S)) = p(B \cup \sigma(W_4) \cup \sigma(S')) = 1$, for some $\sigma(S) = W_2$ and $\sigma(S') = W_5$, thus $\text{dist}(p, \sigma(p)) \geq 4$.

- (b) $\sigma(W_5) = W_5$ and $\sigma(W_2) = W_2$.

Since $\sigma(C) \neq C$, W.L.O.G, assume $\sigma(W_1) \in D$, then $p(B \cup \sigma(W_1) \cup \sigma(W_5)) = 1$.

If $\sigma(W_6) \in D$, then $p(B \cup \sigma(W_2) \cup \sigma(W_6)) = 1$ and $p(B \cup \sigma(S) \cup \sigma(W_5)) = p(B \cup \sigma(S') \cup \sigma(W_2)) = 0$, for some $\sigma(S) = W_1$ and $\sigma(S') = W_6$.

If $\sigma(W_6) = W_6$, then $p(B \cup \sigma(W_1) \cup \sigma(W_6)) = 1$, and $p(B \cup \sigma(S) \cup \sigma(W_5)) = p(B \cup \sigma(S) \cup \sigma(W_6)) = 0$, for some $\sigma(S) = W_1$.

If $\sigma(W_6) = W_1$, then $p(B \cup \sigma(W_2) \cup \sigma(W_6)) = 1$ and $p(B \cup \sigma(W_6) \cup \sigma(W_5)) = 0$.

Thus we always have $dist(p, \sigma(p)) \geq 3$.

(c) $\sigma(W_5) \neq W_5$ and $\sigma(W_2) = W_2$.

Note that $p(B \cup \sigma(W_4) \cup \sigma(W_5)) = 0$ and $p(B \cup \sigma(W_4) \cup \sigma(S)) = 1$ for some $\sigma(S) = W_5$.

If $\sigma(W_5) \in D \cup \{W_1\}$, then $p(B \cup \sigma(W_2) \cup \sigma(W_5)) = 1$.

If $\sigma(W_5) = W_6$ and $\sigma(W_6) \in D \cup \{W_1\}$, then $p(B \cup \sigma(W_2) \cup \sigma(W_6)) = 1$.

If $\sigma(W_5) = W_6$ and $\sigma(W_6) \in W_5$, since $\sigma(C) \neq C$, thus $\sigma(W_1) \in D$ and $p(B \cup \sigma(W_1) \cup \sigma(W_5)) = 1$.

Therefore we always have $dist(p, \sigma(p)) \geq 3$.

(d) $\sigma(W_2) \neq W_2$ and $\sigma(W_5) = W_5$.

Similar to the above one.

4. $\sigma(W_3) = W_4$ and $\sigma(W_4) = W_3$.

Similar to the case where $\sigma(W_3) = W_3$ and $\sigma(W_4) = W_4$.

Case for $k_2 \geq 2$ Similarly, since $|\{\sigma(p)|\sigma(C) = C \text{ and } \sigma(B) = B\}| = O(1)$, we only need to consider the set $\{\sigma(p)|\sigma(C) \neq C \text{ and } \sigma(B) = B\}$, and we exclude each σ case by case:

1. $\sigma(W_3)$ or $\sigma(W_4) \notin \{W_3, W_4\}$.

Assume $\sigma(W_3) \notin \{W_3, W_4\}$, note that for any $S \cap (B \cup W_3) = \emptyset$, $p(B \cup W_3 \cup S) \neq \star$ and there are only two such S s to make $p = 1$. While no matter what $\sigma(W_3)$ is, it's easy to see there are at least five (actually many) such S s to make $p(B \cup \sigma(W_3) \cup \sigma(S)) = 1$, thus $dist(p, \sigma(p)) \geq 3$.

2. $\sigma(W_3), \sigma(W_4) \in \{W_3, W_4\}$

W.L.O.G, assume $\sigma(W_3) = W_3$ and $\sigma(W_4) = W_4$.

(a) $\sigma(W_5) \neq W_5$ and $\sigma(W_2) \neq W_2$.

Now $p(B \cup \sigma(W_3) \cup \sigma(W_2)) = p(B \cup \sigma(W_4) \cup \sigma(W_5)) = 0$ and $p(B \cup \sigma(W_3) \cup \sigma(S)) = p(B \cup \sigma(W_4) \cup \sigma(S')) = 1$, for some $\sigma(S) = W_2$ and $\sigma(S') = W_5$. Therefore, $dist(p, \sigma(p)) \geq 4$.

(b) $\sigma(W_5) = W_5$ or $\sigma(W_2) = W_2$.

Assume $\sigma(W_5) = W_5$, since $\sigma(C) \cap D \neq \emptyset$, there exists some $W \in \{W_1, W_2, W_6\}$ such that $\sigma(W) \cap D \neq \emptyset$.

Moreover, for any $S \subseteq W_3 \cup W_4 \cup W_5$ and $S \notin \{W_3, W_4, W_5\}$ with $|S| = k_2$, we have $\sigma(S) \subseteq W_3 \cup W_4 \cup W_5$ and $\sigma(S) \notin \{W_3, W_4, W_5\}$, thus $p(B \cup W \cup S) = 0 \neq p(B \cup \sigma(W) \cup \sigma(S)) = 1$, and note that there are at least $\binom{6}{2} - 3 = 12$ such S s. Thus, $\text{dist}(p, \sigma(p)) \geq 3$.

□

Proof. (Proof of Theorem 2) We still use minterm functions here.

Let $k = 3l + 1$. Note that in the above construction for $(3l + 1)$ -uniform hypergraph properties, $s_1(f) \leq |p| = O(n^l)$ and $s_0(f) = O(n^{l+1})$. Intuitively, we can pack \sqrt{n} minterms together to get a super minterm, expecting to decrease the number of isomorphisms satisfying the distance condition (i.e., where any of two isomorphisms of p have distance at most 2). Unfortunately, just packing minterms naively doesn't work here, we need some tricks.

Let p be the minterm defining f . p is constructed as follow:

The notions V, B, W_i, C and D are defined the same as in Theorem 1, where we let $k_1 = k_2 = l$. Besides that, let $D_1 = \{v_{6l+1}, v_{6l+2}, \dots, v_{6l+\sqrt{n}}\}$ and $D_2 = D \setminus D_1$.

- For any $S \subseteq C$ of size $2l$ and any $v \in D_1$, $p(B \cup S \cup v) = 0$, except $S = W_i \cup W_{i+1}$ for $i \in [5]$ where $p(B \cup S \cup v) = 1$.
- For any $S \subseteq C$ of size $2l$ and any $v \in D_2$, $p(B \cup S \cup v) = 1$.
- For any S satisfying $l \leq |S \cap C| < 2l$, $|S| = 2l+1$ and $S \cap D_1 \neq \emptyset$, $p(B \cup S) = 1$, except W_3 or $W_4 \subseteq S$ where $p(B \cup S) = 0$.
- All the other variables are \star .

It is not hard to see that $|p| = O(n^{l+1/2})$, thus $s_1(f) \leq C_1(f) \leq |p| = O(n^{l+1/2})$.

Similar to the argument in the proof of Theorem 1, we just need to show the following claim to complete the proof.

Claim 2. *There are only $O(\sqrt{n})$ $\sigma(p)$ s with the same $\pi(B) = \sigma(B)$ satisfying $\text{dist}(\pi(p), \sigma(p)) \leq 2$.*

Proof. By contradiction, suppose there are $C\sqrt{n}$ such $\sigma(p)$ s where C is a sufficient large number, thus there must exist a vertice v such that $\sigma(v) \in D_1$ for at least C such $\sigma(p)$ s, w.l.o.g, assume this set contains p . And we will argue that there are only $O(1)$ such $\sigma(p)$ s satisfying $\text{dist}(\sigma(p), p) \leq 2$, then it's a contradiction, which completes the proof.

Since the vertices in D_1 or D_2 are symmetric, thus $|\{\sigma(p) | \sigma(C) = C \text{ and } \sigma(D_1) = D_1\}| = O(1)$.

If $\sigma(C) = C$ and $\exists v_1 \in D_1, v_2 \in D_2$ satisfying $\sigma(v_1) = v_2$, then $\text{dist}(\sigma(p), p) \geq 3$, since almost all variables which contains v_1, C and B are 0 in p , while all these variable are 1 in $\sigma(p)$.

If $\sigma(C) \neq C$, since $\sigma(v) \in D_1$, then we find that $p(S \cup v) = p'(S)$ where p' is the minterm defined in Theorem 1 for $3l$ -uniform hypergraph properties. Similarly,

$\sigma(p)(S \cup v) = p(\sigma(S) \cup \sigma(v)) = p'(\sigma(S))$. We only consider those S s satisfying $v \notin \sigma(S) \cup S$ and follows the similar proof of Claim 1 in Theorem 1. Finally we can obtain $\text{dist}(p, \sigma(p)) \geq 3$.

□

4 k -Partite k -Uniform Hypergraph Properties and Abelian Groups

In this section, we give the constructions of k -partite k -uniform hypergraph properties first.

Proof. (Proof of Theorem 3)

$\vec{b} =$	$\vec{1}$	$\vec{2}$	$\vec{3}$	$\vec{4}$	$\vec{5}$	$\vec{6}$	$\vec{7}$	\dots
$\vec{a} = \vec{1}$	0	0	0	1	1	1	1	1
$\vec{2}$	0	0	1	1	1	1	1	1
$\vec{3}$	1	0	1	0	0	0	0	0
$\vec{4}$	1	1	0	★	★	★	★	★
$\vec{5}$	1	1	0	★	★	★	★	★
$\vec{6}$	1	1	0	★	★	★	★	★
$\vec{7}$	1	1	0	★	★	★	★	★
\dots	1	1	0	★	★	★	★	★

Table 1: The tabel to illustrate p of k -partite k -uniform hypergraph properties.

The function we use here is also a minterm function. Let k_1 and k_2 be the integers such that $k_1 + 2k_2 = k$ and $k_1, k_2 \leq \lceil k/3 \rceil$. We divide the k partitions into three sets, and each of them is of size k_2 , k_2 and k_1 and indicated by $\vec{a}, \vec{b} \in [n]^{k_2}$ and $\vec{c} \in [n]^{k_1}$ respectively. We use Table 1 to illustrate the minterm p :

- For $\vec{b} = \vec{1}, \vec{2}$ or $\vec{3}$, $p(\vec{1}, \vec{b}, \vec{1}_c) = 0$, otherwise $p(\vec{1}, \vec{b}, \vec{1}_c) = 1$.
- For $\vec{b} = \vec{1}$ or $\vec{2}$, $p(\vec{2}, \vec{b}, \vec{1}_c) = 0$, otherwise $p(\vec{2}, \vec{b}, \vec{1}_c) = 1$.
- For $\vec{b} = \vec{1}$ or $\vec{3}$, $p(\vec{3}, \vec{b}, \vec{1}_c) = 1$, otherwise $p(\vec{3}, \vec{b}, \vec{1}_c) = 0$.
- For $\vec{a} \notin \{\vec{1}, \vec{2}, \vec{3}\}$ and $\vec{b} = \vec{1}$ or $\vec{2}$, $p(\vec{a}, \vec{b}, \vec{1}_c) = 1$.
- For $\vec{a} \notin \{\vec{1}, \vec{2}, \vec{3}\}$ and $\vec{b} = \vec{3}$, $p(\vec{a}, \vec{b}, \vec{1}_c) = 0$.
- Otherwise $p(\vec{a}, \vec{b}, \vec{c}) = \star$.

Here $\vec{1}, \vec{2}$ and $\vec{3}$ can be any three different vectors, W.L.O.G, assume $\vec{1} = (1, \dots, 1, 1)$, $\vec{2} = (1, \dots, 1, 2)$, $\vec{3} = (1, \dots, 1, 3)$ and $\vec{1}_c = (1, \dots, 1, 1)$. It's easy to see $s_1(f) \leq C_1(f) \leq |p| = O(n^{k_2})$. By discussing case by case, it can be verified that for any

p^π there are at most $O(1)$ p^σ s satisfying $\pi(\vec{c}) = \sigma(\vec{c})$ and $dist(p^\pi, p^\sigma) \leq 2$. Thus $s_0(f) = O(n^{k_1})$ since there are at most n^{k_1} choices of \vec{c} . The verify procedure is trivial but tedious, and we omit it here. \square

In the following, we give the proofs of Theorem 4 and Theorem 5.

Proof. (Proof of Theorem 4) We still use minterm functions here. Let $k = 3l + 1$ where $l \geq 1$. We divide the k partitions into four sets of size l, l, l and 1 , and each set is indicated by $\vec{a}, \vec{b}, \vec{c} \in [n]^l$ and $\vec{d} \in [n]$ respectively. The minterm p is constructed as follow:

- For any $\vec{d} \in [\sqrt{n}]$, and any \vec{a} and \vec{b} , $p(\vec{a}, \vec{b}, \vec{1}, \vec{d}) = p'(\vec{a}, \vec{b}, \vec{1}_c)$. Here p' is the partial assignment defined in the proof of Theorem 3.
- For any $\vec{d} \notin [\sqrt{n}]$ and $\vec{a}, \vec{b} \in \{\vec{1}, \vec{2}, \vec{3}\}$, $p(\vec{a}, \vec{b}, \vec{1}, \vec{d}) = 1$.
- Otherwise $p(\vec{a}, \vec{b}, \vec{c}, \vec{d}) = \star$.

It's easy to see $s_1(f) \leq |p| = O(n^{l+1/2})$. It is also not hard to verify that there are at most \sqrt{n} p^σ s with the same $\sigma(\vec{c})$ and satisfying the condition that the distance between any two of them is at most 2, thus $s_0(f) = O(n^{l+1/2})$. \square

Proof. (Proof of Theorem 5) First note that the transitive action of a group G on $[n]$ is equivalent to the action of G by left multiplication on a coset space G/Stab_1 , here Stab_1 is the stabilizer of the element $1 \in [n]$. Since G is an Abelian group, $\text{Stab}_1 = \dots = \text{Stab}_n$, thus $\text{Stab}_1 = \{e\}$. Therefore, the action of G on $[n]$ is equivalent to the action of G by multiplication on itself. So we can relabel the variables (x_1, \dots, x_n) as $(x_{(1, \dots, 1)}, \dots, x_{(m_1, \dots, m_l)})$ to make $(\sigma_1 \otimes \dots \otimes \sigma_l)(x) = (x_{(\sigma_1(1), \dots, \sigma_l(1))}, \dots, x_{(\sigma_1(m_1), \dots, \sigma_l(m_l))})$ for any $\sigma_1 \otimes \dots \otimes \sigma_l \in C_{m_1} \times \dots \times C_{m_l}$.

Let p_m be the minterm of $f : \{0, 1\}^m \rightarrow \{0, 1\}$ defined by Chakraborty in Theorem 3.1 in [13]. We define the minterm p as $p(i_1, \dots, i_l) = \bigoplus_{j=1}^l p_{m_j}(i_j)$. Here $\star \oplus b = \star$, for $b = 0, 1$, or \star . It is easy to see $s_1(f) \leq |p| = \prod_{j=1}^l |p_{m_j}| \leq \gamma n^{1/3}$, where γ is a number only depending on l . Moreover, according to the construction of p_m , it is easy to see that there are at most $\beta n^{1/3}$ $\sigma(p)$ s where the distance between any two of them is at most 2. Here β is another number only depending on l , thus $s_0(f) \leq \beta n^{1/3}$. This completes the proof. \square

5 Lower bounds

In this section, we give the proof of Theorem 6. The proof is similar with Lemma 8 in [11].

Proof. (Proof of Theorem 6) W.O.L.G we assume that for the empty graph \overline{K}_n , $f(\overline{K}_n) = 0$. Since f is non-trivial, there must exist a graph G such that $f(G) = 1$. Let's consider graphs in $f^{-1}(1) = \{G | f(G) = 1\}$ with the minimum number of edges. Define $m = \min\{|E(G)| : f(G) = 1\}$.

We claim that if $m \geq \frac{1}{k+2}n$, then $s(f) \geq \frac{1}{k+2}n$. Let G be a graph in $f^{-1}(1)$ and $|E(G)| = m$. Consider the subfunction f' where $\forall e \notin E(G)$, x_e is restricted to 0, since G has the minimum number of edges, deleting any edges from G will change the values of $f(G)$, therefore, f' is a AND function. Thus, $s(f) \geq s(f') = m \geq \frac{1}{k+2}n$.

In the following we assume $m < \frac{1}{k+2}n$. Again let G be a graph in $f^{-1}(1)$ with $|E(G)| = m$. Let us consider the isolated vertices set I , as

$$\sum_{v \in V} \deg(v) = k|E(G)| < \frac{k}{k+2}n.$$

We have

$$|I| \geq n - \sum_{v \in V} \deg(v) > \frac{2}{k+2}n.$$

Suppose $s(f) < \frac{1}{k+2}n$, we will deduce that there exists another graph with fewer edges and the same value, against the assumption that G has the minimum number of edges in $f^{-1}(1)$, which ends the whole proof.

Pick a vertex u with $\deg(u) = d > 0$. Suppose in the graph G vertex u is adjacent to $(k-1)$ -edges $\{e_1^{(k-1)}, e_2^{(k-1)}, \dots, e_d^{(k-1)}\}$ and $I = \{u_1, u_2, \dots, u_t\}$, where $t = |I|$.

Consider the t -variable Boolean function $g_1: \{0, 1\}^t \rightarrow \{0, 1\}$, where

$$g_1(x_1, \dots, x_t) = f(G + x_1(e_1^{(k-1)}, u_1) + \dots + x_t(e_t^{(k-1)}, u_t)).$$

It is easy to see that g_1 is a symmetric function. We claim that g_1 is a constant function: if not, we have $s(g_1) \geq \frac{1}{2}t$ [10], which implies $s(f) > \frac{1}{k+2}n$ since g_1 is a restriction of f . In particular, $g_1(1, \dots, 1) = g_1(0, \dots, 0)$, i.e. $f(G_1) = f(G)$, where $G_1 = G + \sum_{i=1}^t (e_i^{(k-1)}, u_i)$.

Define $G_i = G_{i-1} + \sum_{j=1}^t (e_j^{(k-1)}, u_j)$ ($i = 2, \dots, d$). Similarly, we can show that

$$f(G) = f(G_1) = \dots = f(G_d).$$

Next we will delete all the edges between $\{u, u_1, \dots, u_t\}$ and $\{e_1^{(k-1)}, e_2^{(k-1)}, \dots, e_d^{(k-1)}\}$ from G_d by reversing the adding edge procedure of $G \rightarrow G_1 \rightarrow \dots \rightarrow G_d$. More precisely, define $H_1 = G_d$; for $i = 2, \dots, d$, define

$$H_i = H_{i-1} - (e_i^{(k-1)}, u) - (e_i^{(k-1)}, u_1) - \dots - (e_i^{(k-1)}, u_t),$$

and

$$h_i(y_0, y_1, \dots, y_t) = f(H_i + y_0(e_i^{(k-1)}, u) + y_1(e_i^{(k-1)}, u_1) + \dots + y_t(e_i^{(k-1)}, u_t)).$$

Similarly, by the fact $s(f) < \frac{1}{k+2}n$ we can show that all the functions h_2, \dots, h_d are constant, which implies $f(H_1) = f(H_2) = \dots = f(H_d)$. So we find another graph H_d with fewer edges than G and $f(H_d) = 1$. \square

6 Conclusion

In this paper, we present a k -uniform hypergraph property with sensitivity complexity $O(n^{\lceil k/3 \rceil})$ for any $k \geq 3$ and we can do better when $k \equiv 1 \pmod{3}$. Besides that, we also investigate the sensitivity complexity of other transitive Boolean functions with certain symmetry. All the functions we constructed in this paper are minterm transitive functions. On the other side, Charkrobati [13] proved that the sensitivity complexity of any minterm transitive Boolean function $f : \{0, 1\}^n \rightarrow \{0, 1\}$ is at least $\Omega(n^{1/3})$. Kulkarni et al. [35] point out that the existence of any transitive function $f : \{0, 1\}^n \rightarrow \{0, 1\}$ with $s(f) = n^\alpha$ where $\alpha < 1/3$ implies a larger than quadratic separation between block sensitivity and sensitivity. We conjecture that the example here is almost tight.

Conjecture 1. *For any constant $k \geq 3$ and for any non-trivial k -hypergraph property f , $s(f) = \Omega(n^{k/3})$, where n is the number of vertices.*

Conjecture 2. *For any $k \geq 3$, there exist a sequence of k -uniform hypergraph properties f with $s(f) = O(n^{k/3})$, where n is the number of vertices.*

A more general question is the following variant of Turan's question proposed by Chakraborty [13]: If f is Boolean function invariant under a transitive group of permutations then is it true that $s(f) = \Omega(n^c)$ for some constant $c > 0$? We conjecture that the inequality holds for $c = 1/3$, which would imply Conjecture 1 and the sensitivity conjecture of transitive functions.

References

- [1] Joshua Biderman, Kevin Cuddy, Ang Li, and Min Jae Song. On the sensitivity of k -uniform hypergraph properties. *CoRR*, abs/1510.00354, 2015.
- [2] László Lovász and Neal E. Young. Lecture notes on evasiveness of graph properties. *CoRR*, cs.CC/0205031, 2002.
- [3] Amit Chakrabarti, Subhash Khot, and Yaoyun Shi. Evasiveness of subgraph containment and related properties. *SIAM J. Comput.*, 31(3):866–875, 2001.
- [4] Raghav Kulkarni. Evasiveness through a circuit lens. In *Innovations in Theoretical Computer Science, ITCS '13, Berkeley, CA, USA, January 9-12, 2013*, pages 139–144, 2013.
- [5] László Babai, Anandam Banerjee, Raghav Kulkarni, and Vipul Naik. Evasiveness and the distribution of prime numbers. In *27th International Symposium on Theoretical Aspects of Computer Science, STACS 2010, March 4-6, 2010, Nancy, France*, pages 71–82, 2010.
- [6] Ronald L. Rivest and Jean Vuillemin. On recognizing graph properties from adjacency matrices. *Theor. Comput. Sci.*, 3(3):371–384, 1976.

- [7] Raghav Kulkarni, Youming Qiao, and Xiaoming Sun. Any monotone property of 3-uniform hypergraphs is weakly evasive. *Theor. Comput. Sci.*, 588:16–23, 2015.
- [8] Timothy Black. Monotone properties of k-uniform hypergraphs are weakly evasive. In *Proceedings of the 2015 Conference on Innovations in Theoretical Computer Science*, ITCS '15, pages 383–391, New York, NY, USA, 2015. ACM.
- [9] Xiaoming Sun. Block sensitivity of weakly symmetric functions. *Theor. Comput. Sci.*, 384(1):87–91, 2007.
- [10] György Turán. The critical complexity of graph properties. *Information Processing Letters*, 18(3):151–153, 1984.
- [11] Xiaoming Sun. An improved lower bound on the sensitivity complexity of graph properties. *Theoretical Computer Science*, 412(29):3524 – 3529, 2011.
- [12] Yihan Gao, Jieming Mao, Xiaoming Sun, and Song Zuo. On the sensitivity complexity of bipartite graph properties. *Theor. Comput. Sci.*, 468:83–91, 2013.
- [13] Sourav Chakraborty. On the sensitivity of cyclically-invariant boolean functions. In *Proceedings of the 20th Annual IEEE Conference on Computational Complexity*, CCC '05, pages 163–167, Washington, DC, USA, 2005. IEEE Computer Society.
- [14] Stephen Cook and Cynthia Dwork. Bounds on the time for parallel ram's to compute simple functions. In *Proceedings of the Fourteenth Annual ACM Symposium on Theory of Computing*, STOC '82, pages 231–233, New York, NY, USA, 1982. ACM.
- [15] Stephen A. Cook, Cynthia Dwork, and Rüdiger Reischuk. Upper and lower time bounds for parallel random access machines without simultaneous writes. *SIAM J. Comput.*, 15(1):87–97, 1986.
- [16] Noam Nisan. Crew prams and decision trees. *SIAM Journal on Computing*, 20(6):999–1007, 1991.
- [17] Harry Buhrman and Ronald De Wolf. Complexity measures and decision tree complexity: a survey. *Theoretical Computer Science*, 288(1):21–43, 2002.
- [18] Noam Nisan and Mario Szegedy. On the degree of boolean functions as real polynomials. In *Proceedings of the Twenty-fourth Annual ACM Symposium on Theory of Computing*, STOC '92, pages 462–467, New York, NY, USA, 1992. ACM.

- [19] Andris Ambainis, Krisjanis Prusis, and Jevgenijs Vihrovs. Sensitivity versus certificate complexity of boolean functions. In *Computer Science - Theory and Applications - 11th International Computer Science Symposium in Russia, CSR 2016, St. Petersburg, Russia, June 9-13, 2016, Proceedings*, pages 16–28, 2016.
- [20] Kun He, Qian Li, and Xiaoming Sun. A tighter relation between sensitivity and certificate complexity. *Manuscript*.
- [21] Andris Ambainis and Xiaoming Sun. New separation between $s(f)$ and $bs(f)$. *Electronic Colloquium on Computational Complexity (ECCC)*, 18:116, 2011.
- [22] Pooya Hatami, Raghav Kulkarni, and Denis Pankratov. Variations on the sensitivity conjecture. *Theory of Computing, Graduate Surveys*, 4:1–27, 2011.
- [23] Meena Boppana. Lattice variant of the sensitivity conjecture. *Electronic Colloquium on Computational Complexity (ECCC)*, 19:89, 2012.
- [24] Andris Ambainis and Krisjanis Prusis. A tight lower bound on certificate complexity in terms of block sensitivity and sensitivity. In *Mathematical Foundations of Computer Science 2014 - 39th International Symposium, MFCS 2014, Budapest, Hungary, August 25-29, 2014. Proceedings, Part II*, pages 33–44, 2014.
- [25] Andris Ambainis, Mohammad Bavarian, Yihan Gao, Jieming Mao, Xiaoming Sun, and Song Zuo. Tighter relations between sensitivity and other complexity measures. In *Automata, Languages, and Programming - 41st International Colloquium, ICALP 2014, Copenhagen, Denmark, July 8-11, 2014, Proceedings, Part I*, pages 101–113, 2014.
- [26] Andris Ambainis and Jevgenijs Vihrovs. Size of sets with small sensitivity: A generalization of simon’s lemma. In *Theory and Applications of Models of Computation - 12th Annual Conference, TAMC 2015, Singapore, May 18-20, 2015, Proceedings*, pages 122–133, 2015.
- [27] Justin Gilmer, Michal Koucký, and Michael E. Saks. A new approach to the sensitivity conjecture. In *Proceedings of the 2015 Conference on Innovations in Theoretical Computer Science, ITCS 2015, Rehovot, Israel, January 11-13, 2015*, pages 247–254, 2015.
- [28] Mario Szegedy. An $o(n^{0.4732})$ upper bound on the complexity of the GKS communication game. *Electronic Colloquium on Computational Complexity (ECCC)*, 22:102, 2015.
- [29] Parikshit Gopalan, Noam Nisan, Rocco A. Servedio, Kunal Talwar, and Avi Wigderson. Smooth boolean functions are easy: Efficient algorithms for low-sensitivity functions. In *Proceedings of the 2016 ACM Conference on In-*

novations in Theoretical Computer Science, Cambridge, MA, USA, January 14-16, 2016, pages 59–70, 2016.

- [30] Parikshit Gopalan, Rocco A. Servedio, and Avi Wigderson. Degree and sensitivity: Tails of two distributions. In *31st Conference on Computational Complexity, CCC 2016, May 29 to June 1, 2016, Tokyo, Japan*, pages 13:1–13:23, 2016.
- [31] Avishay Tal. On the sensitivity conjecture. *Electronic Colloquium on Computational Complexity (ECCC)*, 23:62, 2016.
- [32] Shalev Ben-David. Low-sensitivity functions from unambiguous certificates. *Electronic Colloquium on Computational Complexity (ECCC)*, 23:84, 2016.
- [33] Chengyu Lin and Shengyu Zhang. Sensitivity conjecture and log-rank conjecture for functions with small alternating numbers. *CoRR*, abs/1602.06627, 2016.
- [34] Karthik C. S. and Sébastien Tavenas. On the sensitivity conjecture for disjunctive normal forms. *CoRR*, abs/1607.05189, 2016.
- [35] Raghav Kulkarni and Avishay Tal. On fractional block sensitivity. *Chicago J. Theor. Comput. Sci.*, 2016, 2016.