

Multiplicity and cold-nuclear matter effects from Glauber-Gribov theory at LHC

I.C. Arsene¹, L. Bravina^{1,2}, A.B. Kaidalov³, K. Tywoniuk¹ and E. Zabrodin^{1,2}

¹ Department of Physics, University of Oslo, N-0316 Oslo

² Institute of Nuclear Physics, Moscow State University, RU-119899 Moscow

³ Institute of Theoretical and Experimental Physics, RU-117259 Moscow

E-mail: konrad@fys.uio.no

Abstract. We present predictions for nuclear modification factor in proton-lead collisions at LHC energy 5.5 TeV from Glauber-Gribov theory of nuclear shadowing. We have also made predictions for baseline cold-matter nuclear effects in lead-lead collisions at the same energy.

1. Introduction

The system formed in nucleus-nucleus (AA) collisions at LHC will provide further insight into the dynamics of the deconfined state of nuclear matter. There are also interesting effects anticipated for the initial state of the incoming nuclei related to shadowing of nuclear parton distributions and the space-time picture of the interaction. These should be studied in the more “clean” environment of a proton-nucleus collision. The initial-state effects constitute a baseline for calculation of the density of particles at all rapidities and affect therefore also high- p_{\perp} particle suppression and jet quenching, as well as the total multiplicity.

Both soft and relatively high- p_{\perp} , $p_{\perp} < 10$ GeV/c, particle production in pA at LHC energies probe the low- x gluon distribution of the target nucleus at moderate scales, $Q^2 \sim p_{\perp}^2$, and is therefore mainly influenced by nuclear shadowing. In the Glauber-Gribov theory [1], shadowing at low- x is related to diffractive structure functions of the nucleon, which are studied at HERA. The space-time picture of the interaction is altered from a longitudinally ordered rescattering at low energies, to a coherent interaction of the constituents of the incoming wave-functions at high energy. Shadowing affects both soft and hard processes. Calculation of gluon shadowing was performed in our recent paper [2], where Gribov approach for the calculation of nuclear structure functions was used. The Schwimmer model was used to account for higher-order rescatterings. The gluon diffractive distributions are taken from the most recent experimental parameterizations [3].

2. Particle production at LHC

Shadowing will lead to a suppression both at mid- and forward rapidities in p+Pb collisions at $\sqrt{s} = 5.5$ TeV as seen in Fig. 1. We have plotted the curves for two distinct kinematical scenarios of particle production; one-jet kinematics which may be well motivated for particle production at $p_{\perp} < 2$ GeV/c and two-jet kinematics that apply for high- p_{\perp} particle production. The uncertainty in the curves is due to uncertainty in the parameterization of gluon diffractive distribution functions. Cronin effect is not included in the curves of Fig. 1. We estimate it to be a 10% effect at these energies.

In Fig. 2 we present the suppression due to cold-nuclear effects in Pb+Pb collisions at $\sqrt{s} = 5.5$ TeV as a function of centrality (top) and rapidity (bottom). Also here we present the results for two kinematics.

References

- [1] V.N. Gribov 1969 *Sov. Phys. JETP* **29** 483
- [2] I.C. Arsene, L. Bravina, A. Kaidalov, K. Tywoniuk and E. Zabrodin, [hep-ph/0705.1596](https://arxiv.org/abs/hep-ph/0705.1596)
- [3] H1 Collaboration, A. Aktas *et al.* 2006 *Eur. Phys. J. C* **48** 715; *ibid.* **48** 749




Figure 1. Shadowing as a function of transverse momentum for p+Pb collisions at $\sqrt{s} = 5.5$ TeV.





Figure 2. Shadowing as a function of centrality (top) and rapidity (bottom) for Pb+Pb collisions at $\sqrt{s} = 5.5$ TeV.