

## SHNIRELMAN'S THEOREM APPLICATIONS

FELIX SIDOKHINE

ABSTRACT. Shnirelman's theorem is applied to solving Diophantine equations, and also discussing of the problems of a representation of Gaussian integers by a sum of odd Gaussian primes.

## 1. INTRODUCTION

In our note we study the systems of Diophantine equations and connect this problem to Shnirelman's theorem: "Every integer  $n > 7$  is a sum at most  $s_0$  (Shnirelman's constant) odd primes" [1]. The problem of the representation of Gaussian integers by a sum of odd Gaussian primes is explored from an angle of Shnirelman's theorem.

## 2. DIOPHANTINE EQUATIONS AND SHNIRELMAN'S THEOREM

**Theorem 1.** *The system of diophantine equations:*

$$\begin{cases} x_{11} + x_{12} + x_{13} + x_{14} = a \\ x_{21} + x_{22} + x_{23} + x_{24} = b \\ \forall s \ x_{1s} + x_{2s} = p_s \text{ where } p_s \text{ is an odd prime} \end{cases}$$

has a solution  $(x_{11}, \dots, x_{24})$  where  $x_{ij} \in \mathbb{Z}_+^1$  if  $a > 10$ ,  $a \geq b > 0$  and  $a + b \equiv 0 \pmod{2}$ .

*Proof.* The proof uses induction. Let the theorem be true up to some  $n - 2 = a + b$ . Let the theorem be false for  $n = a' + b'$ . By Shnirelman's theorem with  $s_0 = 4$ , there exist  $p \geq q \geq r \geq l \in \pi_{\mathbb{Z}}^2$  such that  $n = a' + b' = p + q + r + l$ . Let us consider 4 cases:

- Case 1:  $b' \leq l$ , then we have a solution: 
$$\begin{cases} p + q + r + (l - b') = a' \\ 0 + 0 + 0 + b' = b' \end{cases}$$
- Case 2:  $l < b' \leq r + l$ , then we have a solution: 
$$\begin{cases} p + q + (r + l - b') + 0 = a' \\ 0 + 0 + (b' - l) + l = b' \end{cases}$$
- Case 3:  $r + l < b' \leq q + r + l$ , then we have a solution: 
$$\begin{cases} p + (q + r + l - b') + 0 + 0 = a' \\ 0 + (b' - r - l) + r + l = b' \end{cases}$$

---

Date: January 3, 2020.

<sup>1</sup> $\mathbb{Z}_+ = \{n \in \mathbb{Z} | n \geq 0\}$

<sup>2</sup> $\pi_{\mathbb{Z}} = \{n \in \mathbb{Z} | n \text{ is prime}\}$

- Case 4:  $q+r+l < b' < p+q+r+l$ , then we have a solution: 
$$\begin{cases} (p+q+r+l-b') + 0 + 0 + 0 = a' \\ (b' - q - r - l) + q + r + l = b' \end{cases}$$

From this, it follows that the theorem is true. □

**Theorem 2.** *Given a system of diophantine equations where  $a > 6$  and  $a \geq b > 0$ :*

$$\begin{cases} \sum_{i=1}^k x_{1i} = a \\ \sum_{j=1}^k x_{2j} = b \\ \forall s \ x_{1s} + x_{2s} = p_s \text{ where } p_s \text{ is an odd prime} \end{cases}$$

there exists such  $k$ , where  $k \leq k_0$  (constant), that the system has a solution  $(x_{11}, \dots, x_{2n})$  where  $x_{ij} \in \mathbb{Z}_+$ .

*Proof.* The proof mimics the one used for theorem 1. □

Based on these two theorems, we can propose the following conjecture which will be useful to use when working with Gaussian integers.

**Conjecture 1.** *Given the system of equations where  $a > 6$  and  $a \geq b > 0$ :*

$$\begin{cases} \sum_{i=1}^k x_{1i} = a \\ \sum_{j=1}^k x_{2j} = b \\ \forall s \ x_{1s}^2 + x_{2s}^2 = t_s \text{ where } t_s \text{ is either a prime } p_s \equiv 1 \pmod{4} \text{ or } t_s = p_s^2 \text{ where } p_s \equiv 3 \pmod{4}. \end{cases}$$

there exists such  $k$ , where  $k \leq k_1$  (constant), that the system has a solution  $(x_{11}, \dots, x_{2n})$  where  $x_{ij} \in \mathbb{Z}_+$ .

Let us consider  $z \in \mathbb{Z}[i]$  and let  $z = a + ib$ . If the above conjecture is true, then it implies that  $z$  is a sum of at most  $k_1$  odd Gaussian primes with non-negative real and imaginary parts [2].

### 3. GAUSSIAN INTEGERS, ODD GAUSSIAN PRIMES AND SHNIRELMAN'S THEOREM

Let us introduce the notion of “odd” and “even” Gaussian integers, where  $z$  is odd if  $z \equiv 1 \pmod{1+i}$  and even if  $z \equiv 0 \pmod{1+i}$ .

Our global experiment was to see whenever any element from the set  $\Gamma_G = \{z \in \mathbb{Z}[i] \mid \operatorname{Re}(z) > 0 \wedge -\operatorname{Re}(z) < \operatorname{Im} z \leq \operatorname{Re}(z)\}$  could be represented as a sum of at most  $k_0$  elements from the set  $\Gamma_\pi = \{z \in \pi_{\mathbb{Z}[i]} \mid \operatorname{Re}(z) > 0 \wedge -\operatorname{Re}(z) < \operatorname{Im}(z) \leq \operatorname{Re}(z)\}$ . Some of our computations and results are presented in appendix.

Our experimental work has allowed us to observe some interesting facts about Gaussian integers, notably given  $z \in \Gamma_G$ , and either  $\operatorname{Im}(z) = \operatorname{Re}(z) - 1$  or  $\operatorname{Im}(z) = \operatorname{Re}(z)$ ,  $z$  cannot be represented as a sum of odd primes from  $\Gamma_\pi$ . This has led us to consider a different choice for our set  $\Gamma_G = \{z \in \mathbb{Z}[i] \mid \operatorname{Re}(z) > 0 \wedge \operatorname{Im}(z) \geq 0\}$  and the following set  $K_\pi = \{z \in \pi_{\mathbb{Z}[i]} \mid \operatorname{Re}(z) \geq 0 \wedge \operatorname{Im}(z) \geq 0\}$ . It is worth noting that in both cases  $\Gamma_G$  is a maximal set of non-associated Gaussian integers.

Studying the results from [2] and our experimental data we have formulated the following conjecture:

**Conjecture 2.** *Given  $A = \{z \in \mathbb{Z}[i] \mid \operatorname{Re}(z) > 0 \wedge \operatorname{Im}(z) > 0\}$  then for all  $z$  in  $A$ , such that  $\max(\operatorname{Re}(z), \operatorname{Im}(z)) \geq 7$ ,  $z$  is a sum of at most 3 odd primes from the set  $K_\pi$ .*

In our study of this conjecture, we have formulated the following proposition:

**Proposition 1.** *Given that for any  $z_0 \in A$  and some integer  $k \geq 2$ , such that  $z_0 \equiv k \pmod{1+i}$  and  $\max(\operatorname{Re}(z_0), \operatorname{Im}(z_0)) \geq c_0 > 0$ , the equation  $\sum_{i=1}^k x_i = z_0$  has a solution  $(x_1, \dots, x_k)$  where  $x_i \in K_\pi$ . Then for any  $w_0 \in A$  where  $w_0 \equiv k+1 \pmod{1+i}$  and  $\max(\operatorname{Re}(w_0), \operatorname{Im}(w_0)) \geq c_0 + 3$ , the equation  $\sum_{i=1}^{k+1} x_i = w_0$  has a solution  $(x_1, \dots, x_{k+1})$  where  $x_i \in K_\pi$ .*

*Proof.* The proof is by induction. Let the proposition be true for some  $k_0$ . Let it be false for  $k_0 + 1$ , therefore there exists such  $w$  that  $w \equiv k_0 + 1 \pmod{1+i}$  and  $\max(\operatorname{Re}(w), \operatorname{Im}(w)) \geq c_0 + 3$  for which the proposition fails. Let  $\operatorname{Im}(w) \geq c_0 + 3$ , then  $w - 3i \equiv k_0 \pmod{1+i}$  and  $\max(\operatorname{Re}(w - 3i), \operatorname{Im}(w - 3i)) \geq c_0$  contradicting the inductive hypothesis. The case  $\operatorname{Re}(w) \geq c_0 + 3$  is analogous.  $\square$

If we assume that every odd Gaussian integer  $z$  where  $\operatorname{Re}(z) > 0$ ,  $\operatorname{Im}(z) > 0$  and  $\max(\operatorname{Re}(z), \operatorname{Im}(z)) \geq c_0$  is a sum of 3 odd Gaussian primes with non-negative real and imaginary parts, then we can use proposition 1 to get the following theorem:

**Theorem 3.** *Any element  $a \in \{z \in \mathbb{Z}[i] \mid \operatorname{Re}(z) > 0 \wedge \operatorname{Im}(z) > 0 \wedge \max(\operatorname{Re}(z), \operatorname{Im}(z)) > c_1\}$  is a sum of at most 4 odd Gaussian primes belonging to the set  $K_\pi = \{z \in \pi_{\mathbb{Z}[i]} \mid \operatorname{Re}(z) \geq 0 \wedge \operatorname{Im}(z) \geq 0\}$ .*

If we consider  $\mathbb{Z}$  as a subset of  $\mathbb{Z}[i]$ , there are even elements of  $\mathbb{Z}$  which can not be represented as a sum of 2 elements from the set  $\pi_{\mathbb{Z}} \cap \pi_{\mathbb{Z}[i]}$ . However, if one of the hypotheses below is true then the elements of  $\mathbb{Z}$  as a subset of  $\mathbb{Z}[i]$  could be represented as a sum of at most  $s_0$  primes from  $\pi_{\mathbb{Z}} \cap \pi_{\mathbb{Z}[i]}$ .

- Hypothesis 1: There exists  $c_0 > 0$  such that every integer  $n \geq c_0$ ,  $n \equiv 2 \pmod{4}$  is a sum of 2 primes where  $p_i \equiv 3 \pmod{4}$
- Hypothesis 2: There exists  $c_0 > 0$  such that every integer  $n \geq c_0$ ,  $n \equiv 1 \pmod{4}$  is a sum of 3 primes where  $p_i \equiv 3 \pmod{4}$
- Hypothesis 3: There exists  $c_0 > 0$  such that every integer  $n \geq c_0$ ,  $n \equiv 0 \pmod{4}$  is a sum of 4 primes where  $p_i \equiv 3 \pmod{4}$
- Hypothesis 4: There exists  $c_0 > 0$  such that every integer  $n \geq c_0$ ,  $n \equiv 3 \pmod{4}$  is a sum of 5 primes where  $p_i \equiv 3 \pmod{4}$

Indeed if at least one of the above hypotheses is true, then the following theorem is also true:

**Theorem 4.** *For all  $n \in \mathbb{Z}$  where  $n \geq c_1$  and  $c_1$  is a constant, there exists  $m$  such that  $\sum_{i=1}^m x_i = n$  has a solution  $(p_1, \dots, p_m)$  where  $\forall i p_i \equiv 3 \pmod{4}$ , and  $p_i$  is prime and  $m$  takes the values ranging from 2 to  $s_1$  (constant).*

*Proof.* Let hypothesis 2 be true and consider the following 3 cases:

- $\forall n$  where  $n \equiv 0 \pmod{4} \wedge n \geq c_0 + 3$  is a sum of four odd primes. Since  $n - 3 \equiv 1 \pmod{4}$  and  $n - 3 \geq c_0$  the proof follows hypothesis 2.

- $\forall n$  where  $n \equiv 3 \pmod{4} \wedge n \geq c_0 + 6$  is a sum of five odd primes. This is true due to the previous case.
- $\forall n$  where  $n \equiv 2 \pmod{4} \wedge n \geq c_0 + 9$  is a sum of six odd primes. This is true due to the previous case.

From these cases, we can conclude that  $c_1 = c_0 + 9$  and  $s_1 = 6$ . □

Using conjecture 2 and theorem 4 we can formulate the following hypothesis:

**Conjecture 3.** *For all  $z \in \Gamma_G$ , where  $\max(\operatorname{Re}(z), \operatorname{Im}(z)) \geq c_0$ ,  $z$  is a sum of at most  $k_0$  odd primes from the set  $K_\pi = \{z \in \pi_{\mathbb{Z}[i]} \mid \operatorname{Re}(z) \geq 0 \wedge \operatorname{Im}(z) \geq 0\}$ , where  $k_0$  is a constant.*

Based on our experimental evidence, we can also propose the following conjecture:

**Conjecture 4.** *For all  $z \in \Gamma_G$ , where  $\max(\operatorname{Re}(z), \operatorname{Im}(z)) \geq 6$ ,  $z$  is a sum of at most 3 odd Gaussian primes from the set  $S_\pi = \{z \in \pi_{\mathbb{Z}[i]} \mid \operatorname{Re}(z) \geq 0 \wedge -\operatorname{Re}(z) < \operatorname{Im}(z)\}$  (i.e.  $z = \sum_{l=1}^k p_l$  where  $p_l \in S_\pi$  is an odd Gaussian prime and  $\|p_l\| < \|z\|$ ).*

#### 4. DISCUSSION AND CONCLUSION

In the present note we have constructed a solution of a system of Diophantine equations of the first degree in several variables over a set of non-negative integers by the Shnirelman theorem. We have demonstrated this approach on Gaussian integers. Based on experimental data on the representation of Gaussian integers by a sum of Gaussian primes we have shown that there is an infinite number of elements of a maximal set of non-associated Gaussian integers  $\Gamma_G$  which cannot be represented as a sum of two or three Gaussian primes belonging to a set  $K_\pi$  ( $\Gamma_G \subset K_\pi \subset \mathbb{Z}[i]$ ). In this case we have formulated the Goldbach conjecture for  $\mathbb{Z}[i]$  as some analogy of the Shnirelman theorem. However, according to our experimental evidence we expect that any Gaussian integer of  $\Gamma_G$ , excepting a finite number, can represent by a sum at most three odd Gaussian primes belonging to a set  $S_\pi$  ( $\Gamma_G \subset K_\pi \subset S_\pi \subset \mathbb{Z}[i]$ ) where  $S_\pi$  is a minimal subset of  $\mathbb{Z}[i]$  to satisfy such conditions.

#### REFERENCES

- [1] O. Ramaré, "Prime numbers: Emergence and victories of bilinear forms decomposition," *European Math. Soc. Newsletter*, pp. 18–28, December 2013.
- [2] O. Knill. Some experiments in number theory. [Online]. Available: <http://arxiv.org/abs/1606.05971>

## 5. APPENDIX

Table 1: Odd Gaussian integers as a linear combination of odd Gaussian primes where  $\Gamma = \{z \in \mathbb{Z}[i] \mid 0 \leq \text{Im}(z) \leq \text{Re}(z)\}$  and  $\Gamma_\pi = \{z \in \pi_{\mathbb{Z}[i]} \mid \text{Re}(z) > 0 \wedge -\text{Re}(z) < \text{Im}(z) \leq \text{Re}(z)\}$

$z \in \Gamma$	$p, q, r \in \Gamma_\pi; \max(\ p\ , \ q\ , \ r\ ) < \ z\ $			Representation
$z = 9$	$p = 3$	$q = 3$	$r = 3$	$z = p + q + r$
$z = 13$	$p = 7$	$q = 3$	$r = 3$	$z = p + q + r$
$z = 17$	$p = 11$	$q = 3$	$r = 3$	$z = p + q + r$
$z = 21$	$p = 11$	$q = 7$	$r = 3$	$z = p + q + r$
$z = 25$	$p = 11$	$q = 11$	$r = 3$	$z = p + q + r$
$z = 29$	$p = 19$	$q = 7$	$r = 3$	$z = p + q + r$
$z = 33$	$p = 19$	$q = 7$	$r = 7$	$z = p + q + r$
$z = 37$	$p = 19$	$q = 11$	$r = 7$	$z = p + q + r$

$z = 7$	$p = 3 + 2i$	$q = 2 - i$	$r = 2 - i$	$z = p + q + r$
$z = 11$	$p = 7$	$q = 2 + i$	$r = 2 - i$	$z = p + q + r$
$z = 15$	$p = 8 + 3i$	$q = 5 - 2i$	$r = 2 - i$	$z = p + q + r$
$z = 19$	$p = 11$	$q = 4 + i$	$r = 4 - i$	$z = p + q + r$
$z = 23$	$p = 13 + 2i$	$q = 7$	$r = 3 - 2i$	$z = p + q + r$
$z = 27$	$p = 13 + 2i$	$q = 11$	$r = 3 - 2i$	$z = p + q + r$
$z = 31$	$p = 23$	$q = 4 + i$	$r = 4 - i$	$z = p + q + r$
$z = 35$	$p = 23$	$q = 6 + i$	$r = 6 - i$	$z = p + q + r$

$z = 6 + i$	$p = 2 + i$	$q = 2 + i$	$r = 2 - i$	$z = p + q + r$
$z = 6 + 3i$	$p = 2 + i$	$q = 2 + i$	$r = 2 + i$	$z = p + q + r$
$z = 7 + 2i$	$p = 3$	$q = 2 + i$	$r = 2 + i$	$z = p + q + r$
$z = 7 + 4i$	$p = 3 + 2i$	$q = 2 + i$	$r = 2 + i$	$z = p + q + r$
$z = 8 + i$	$p = 2 + i$	$q = 3$	$r = 3$	$z = p + q + r$
$z = 8 + 3i$	$p = 3 + 2i$	$q = 3$	$r = 2 + i$	$z = p + q + r$
$z = 8 + 5i$	$p = 3 + 2i$	$q = 3 + 2i$	$r = 2 + i$	$z = p + q + r$
$z = 9 + 2i$	$p = 3 + 2i$	$q = 3$	$r = 3$	$z = p + q + r$
$z = 9 + 4i$	$p = 5 + 2i$	$q = 2 + i$	$r = 2 + i$	$z = p + q + r$
$z = 9 + 6i$	$p = 5 + 4i$	$q = 2 + i$	$r = 2 + i$	$z = p + q + r$
$z = 19 + 16i$	$p = 10 + 9i$	$q = 6 + 5i$	$r = 3 + 2i$	$z = p + q + r$
$z = 27 + 24i$	$p = 23 + 22i$	$q = 2 + i$	$r = 2 + i$	$z = p + q + r$
$z = 39 + 36i$	$p = 35 + 34i$	$q = 2 + i$	$r = 2 + i$	$z = p + q + r$
$z = 48 + 45i$	$p = 25 + 24i$	$q = 20 + 19i$	$r = 3 + 2i$	$z = p + q + r$
$z = 50 + i$	$p = 23$	$q = 20 + i$	$r = 7$	$z = p + q + r$
$z = 50 + 3i$	$p = 40 + i$	$q = 6 + i$	$r = 4 + i$	$z = p + q + r$
$z = 50 + 7i$	$p = 40 + i$	$q = 5 + 4i$	$r = 5 + 2i$	$z = p + q + r$
$z = 50 + 11i$	$p = 36 + 5i$	$q = 9 + 4i$	$r = 5 + 2i$	$z = p + q + r$
$z = 50 + 15i$	$p = 35 + 8i$	$q = 10 + 3i$	$r = 5 + 4i$	$z = p + q + r$
$z = 50 + 19i$	$p = 35 + 16i$	$q = 10 + i$	$r = 5 + 2i$	$z = p + q + r$

$z = 50 + 25i$	$p = 35 + 18i$	$q = 10 + 3i$	$r = 5 + 4i$	$z = p + q + r$
$z = 50 + 41i$	$p = 45 + 38i$	$q = 3 + 2i$	$r = 2 + i$	$z = p + q + r$
$z = 50 + 43i$	$p = 35 + 34i$	$q = 10 + 7i$	$r = 5 + 2i$	$z = p + q + r$
$z = 50 + 45i$	$p = 25 + 24i$	$q = 20 + 19i$	$r = 5 + 2i$	$z = p + q + r$
$z = 50 + 47i$	$p = 25 + 24i$	$q = 20 + 19i$	$r = 5 + 4i$	$z = p + q + r$

$z = 6 + 5i$	$p = 3 + 2i$	$q = 2 + i$	$r = 2 - i$	$z = p + q + ir$
$z = 7 + 6i$	$p = 3 + 2i$	$q = 3 + 2i$	$r = 2 - i$	$z = p + q + ir$
$z = 8 + 7i$	$p = 5 + 4i$	$q = 2 + i$	$r = 2 - i$	$z = p + q + ir$
$z = 9 + 8i$	$p = 5 + 4i$	$q = 3 + 2i$	$r = 2 - i$	$z = p + q + ir$
$x = 11 + 10i$	$p = 8 + 3i$	$q = 2 + i$	$r = 6 - i$	$z = p + q + ir$
$z = 19 + 18i$	$p = 10 + 9i$	$q = 8 + 5i$	$r = 4 - i$	$z = p + q + ir$
$z = 27 + 26i$	$p = 19 + 16i$	$q = 5 + 2i$	$r = 8 - 3i$	$z = p + q + ir$
$z = 39 + 38i$	$p = 35 + 34i$	$q = 3 + 2i$	$r = 2 - i$	$z = p + q + ir$
$z = 48 + 47i$	$p = 23 + 22i$	$q = 20 + 19i$	$r = 6 - 5i$	$z = p + q + ir$
$z = 49 + 48i$	$p = 25 + 24i$	$q = 20 + 19i$	$r = 5 - 4i$	$z = p + q + ir$
$z = 50 + 49i$	$p = 25 + 24i$	$q = 20 + 19i$	$r = 6 - 5i$	$z = p + q + ir$

Table 2: Even Gaussian integers as a linear combination of odd Gaussian primes where  $\Gamma = \{z \in \mathbb{Z}[i] \mid 0 \leq \text{Im}(z) \leq \text{Re}(z)\}$  and  $\Gamma_\pi = \{z \in \pi\mathbb{Z}[i] \mid \text{Re}(z) > 0 \wedge -\text{Re}(z) < \text{Im}(z) \leq \text{Re}(z)\}$

$z \in \Gamma$	$p, q \in \Gamma_\pi; \max(\ p\ , \ q\ ) < \ z\ $		Representation
$z = 6$	$p = 3$	$q = 3$	$z = p + q$
$z = 10$	$p = 7$	$q = 3$	$z = p + q$
$z = 14$	$p = 11$	$q = 3$	$z = p + q$
$z = 18$	$p = 11$	$q = 7$	$z = p + q$
$z = 30$	$p = 19$	$q = 11$	$z = p + q$
$z = 42$	$p = 23$	$q = 19$	$z = p + q$

$z = 8$	$p = 5 + 2i$	$q = 3 - 2i$	$z = p + q$
$z = 12$	$p = 6 + i$	$q = 6 - i$	$z = p + q$
$z = 16$	$p = 13 + 2i$	$q = 3 - 2i$	$z = p + q$
$z = 20$	$p = 10 + i$	$q = 10 - i$	$z = p + q$
$z = 24$	$p = 12 + 7i$	$q = 12 - 7i$	$z = p + q$
$z = 28$	$p = 15 + 2i$	$q = 13 - 2i$	$z = p + q$
$z = 32$	$p = 17 + 2i$	$q = 15 - 2i$	$z = p + q$
$z = 36$	$p = 21 + 4i$	$q = 15 - 4i$	$z = p + q$
$z = 44$	$p = 27 + 2i$	$q = 17 - 2i$	$z = p + q$

$z = 6 + 2i$	$p = 3 + 2i$	$q = 3$	$z = p + q$
$z = 6 + 4i$	$p = 3 + 2i$	$q = 3 + 2i$	$z = p + q$
$z = 7 + i$	$p = 4 + i$	$q = 3$	$z = p + q$

$z = 7 + 3i$	$p = 5 + 2i$	$q = 2 + i$	$z = p + q$
$z = 7 + 5i$	$p = 5 + 4i$	$q = 2 + i$	$z = p + q$
$z = 9 + 7i$	$p = 6 + 5i$	$q = 3 + 2i$	$z = p + q$
$z = 19 + 17i$	$p = 13 + 12i$	$q = 6 + 5i$	$z = p + q$
$z = 27 + 25i$	$p = 25 + 24i$	$q = 2 + i$	$z = p + q$
$z = 36 + 2i$	$p = 26 + i$	$q = 10 + i$	$z = p + q$
$z = 39 + 37i$	$p = 36 + 35i$	$q = 3 + 2i$	$z = p + q$
$z = 48 + 46i$	$p = 25 + 24i$	$q = 23 + 22i$	$z = p + q$
$z = 49 + 47i$	$p = 26 + 25i$	$q = 23 + 22i$	$z = p + q$
$z = 50 + 48i$	$p = 25 + 24i$	$q = 25 + 24i$	$z = p + q$
$z = 51 + i$	$p = 47$	$q = 4 + i$	$z = p + q$
$z = 51 + 3i$	$p = 45 + 2i$	$q = 6 + i$	$z = p + q$
$z = 51 + 7i$	$p = 45 + 2i$	$q = 6 + 5i$	$z = p + q$
$z = 51 + 9i$	$p = 47 + 8i$	$q = 4 + i$	$z = p + q$
$z = 51 + 25i$	$p = 48 + 23i$	$q = 3 + 2i$	$z = p + q$
$z = 51 + 31i$	$p = 49 + 30i$	$q = 2 + i$	$z = p + q$
$z = 51 + 49i$	$p = 48 + 47i$	$q = 3 + 2i$	$z = p + q$

$z = 6 + 6i$	$p = 5 + 4i$	$q = 2 - i$	$z = p + iq$
$z = 7 + 7i$	$p = 5 + 4i$	$q = 3 - 2i$	$z = p + iq$
$z = 8 + 8i$	$p = 7 + 2i$	$q = 6 - i$	$z = p + iq$
$z = 9 + 9i$	$p = 5 + 4i$	$q = 5 - 4i$	$z = p + iq$
$z = 10 + 10i$	$p = 6 + 5i$	$q = 5 - 4i$	$z = p + iq$
$z = 11 + 11i$	$p = 6 + 5i$	$q = 6 - 5i$	$z = p + iq$
$z = 12 + 12i$	$p = 11 + 6i$	$q = 6 - i$	$z = p + iq$
$z = 16 + 16i$	$p = 14 + i$	$q = 15 - 2i$	$z = p + iq$
$z = 19 + 19i$	$p = 18 + 17i$	$q = 2 - i$	$z = p + iq$
$z = 27 + 27i$	$p = 25 + 24i$	$q = 3 - 2i$	$z = p + iq$
$z = 39 + 39i$	$p = 35 + 34i$	$q = 5 - 4i$	$z = p + iq$
$z = 48 + 48i$	$p = 46 + 41i$	$q = 7 - 2i$	$z = p + iq$
$z = 49 + 49i$	$p = 25 + 24i$	$q = 25 - 24i$	$z = p + iq$
$z = 50 + 50i$	$p = 48 + 47i$	$q = 3 - 2i$	$z = p + iq$
$z = 51 + 51i$	$p = 26 + 25i$	$q = 26 - 25i$	$z = p + iq$