An asymmetric multiparameter CCR flow

S. Sundar

January 3, 2020

Abstract

In this note, we exhibit an example of a multiparameter CCR flow which is not cocycle conjugate to its opposite. This is in sharp contrast to the one parameter situation.

AMS Classification No.: Primary 46L55; Secondary 46L99.

Keywords: Decomposable product systems, opposite of a CCR flow.

1 Opposite of an E_0 -semigroup

Let $P \subset \mathbb{R}^d$ be a closed convex cone. We assume that $d \geq 2$, P spans \mathbb{R}^d and P contains no line, i.e. $P \cap -P = \{0\}$. We denote the interior of P by Ω . For $x, y \in \mathbb{R}^d$, we write $x \leq y$ (x < y) if $y - x \in P$ $(y - x \in \Omega)$. Let $\alpha := \{\alpha_x\}_{x \in P}$ be an E_0 -semigroup over P on $B(\mathcal{H})$ where \mathcal{H} is an infinite dimensional separable Hilbert space. For $x \in P$, let

$$E(x) := \{ T \in B(\mathcal{H}) : \alpha_x(A)T = TA \text{ for all } A \in B(\mathcal{H}) \}.$$

For $x \in P$, E(x) is a separable Hilbert space where the inner product is given by $\langle T|S\rangle = S^*T$. The disjoint union of Hilbert spaces $\mathcal{E} := \coprod_{x \in \Omega} E(x)$ has a structure of a product system and is called the product system associated with α . It is indeed a cocycle conjugacy invariant. For more on product systems and E_0 -semigroups in the multiparameter context, we refer the reader to [5].

Keep the foregoing notation. Let

$$\mathcal{E}^{op} := \{ (x, T) \in \Omega \times B(\mathcal{H}) : x \in \Omega, T \in E(x) \}.$$

Define a semigroup multiplication on \mathcal{E}^{op} by the following formula:

$$(x,T).(y,S) = (x+y,ST)$$
 (1.1)

for $(x, T), (y, S) \in \mathcal{E}^{op}$. Then \mathcal{E}^{op} is an abstract product system over Ω (in the sense of Definition 2.1 of [5]). Arveson's bijection between product systems and E_0 -semigroups, established in [5] for the case of a cone, ensures that there exists an E_0 -semigroup denoted $\alpha^{op} := \{\alpha_x^{op}\}_{x\in P}$, which is unique up to cocycle conjugacy, such that the product system associated to α^{op} is isomorphic to \mathcal{E}^{op} . The E_0 -semigroup α^{op} is called the opposite of α .

A natural question that arises in this context is the following. Are the E_0 -semigroups α and α^{op} cocycle conjugate? If α is cocycle conjugate to α^{op} , we call α symmetric otherwise we call α asymmetric. In the one parameter context, we have the following.

- (1) One parameter CCR flows are symmetric. This follows from Arveson's classification of type I E_0 -semigroups and the index computation.
- (2) Tsirelson in his remarkable papers [8] and [9] constructed examples of type II_0 E_0 -semigroups which are asymmetric by probabilistic means.

It is natural to ask whether (1) stays true in the multiparameter context. More precisely, suppose $d \geq 2$ and V is a pure isometric representation of P on a Hilbert space \mathcal{H} . Let α^V be the CCR flow associated to V acting on $B(\Gamma(\mathcal{H}))$ where $\Gamma(\mathcal{H})$ is the symmetric Fock space of \mathcal{H} . Is α^V symmetric? We show that for the left regular representation of P on $L^2(P)$, the associated CCR flow is asymmetric.

2 Decomposable product systems

Following Arveson, the author in [7] has defined the notion of a decomposable product system. Let us review the definitions. Let $\alpha := \{\alpha_x\}_{x \in P}$ be an E_0 -semigroup on $B(\mathcal{H})$ and let $E := \{E(x)\}_{x \in P}$ be the associated product system. Suppose $x \in P$ and $u \in E(x)$ is a non-zero vector. We say that u is decomposable if given $y \leq x$ with $y \in P$, there exists $v \in E(y)$ and $w \in E(x - y)$ such that u = vw. The set of decomposable vectors is denoted by D(x). We say that α is decomposable if

- (1) for $x, y \in P$, D(x)D(y) = D(x + y), and
- (2) for $x \in P$, D(x) is total in E(x).

Proposition 2.1 The opposite of a decomposable E_0 -semigroup is decomposable.

Remark 2.2 The subtle point that we wish to stress is that apriori the product rule in the opposite product system given by Eq. 1.1 holds only over Ω . However to prove Prop. 2.1, we need the validity of the product rule over the whole semigroup P which is assured by the next lemma.

Let us fix a few notation. Let $\alpha := \{\alpha_x\}_{x \in P}$ be a decomposable E_0 -semigroup acting on $B(\mathcal{H})$. The product system of α is denoted by $E := \{E(x)\}_{x \in P}$. We denote the opposite of α by $\beta := \{\beta_x\}_{x \in P}$. Suppose that β acts on $B(\widetilde{\mathcal{H}})$. Denote the product system of β by $F := \{F(x)\}_{x \in P}$.

Lemma 2.3 For every $x \in P$, there exists a unitary $\widetilde{\theta}_x : E(x) \to F(x)$ such that for $x, y \in P$, $T \in E(x)$ and $S \in E(y)$,

$$\widetilde{\theta}_{x+y}(ST) = \widetilde{\theta}_x(T)\widetilde{\theta}_y(S).$$

Proof. From the definition of β , it follows that for $x \in \Omega$, there exists a unitary operator $\theta_x : E(x) \to F(x)$ such that for $x, y \in \Omega$, $T \in E(x)$ and $S \in E(y)$,

$$\theta_{x+y}(ST) = \theta_x(T)\theta_y(S).$$

Fix an element $a \in \Omega$. Let $x \in P$ and $T \in E(x)$ be given. We claim that there exists a unique bounded linear operator on $\widetilde{\mathcal{H}}$, which we denote by $\widetilde{\theta}_x(T)$, such that for $S \in E(a)$ and $\xi \in \widetilde{\mathcal{H}}$,

$$\widetilde{\theta}_x(T)\theta_a(S)\xi = \theta_{a+x}(ST)\xi.$$

For any $b \in \Omega$, the map $E(b) \otimes \widetilde{\mathcal{H}} \ni S \otimes \xi \to \theta_b(S)\xi \in \widetilde{\mathcal{H}}$ is a unitary operator. This way, we can identify $\widetilde{\mathcal{H}}$ with $E(b) \otimes \widetilde{\mathcal{H}}$ for every $b \in \Omega$. Right multiplication by T induces a bounded linear operator from $E(a) \to E(a+x)$ of norm ||T||. Tensor with identity to obtain the desired operator $\widetilde{\theta}_x(T)$ from $\widetilde{\mathcal{H}} \equiv E(a) \otimes \widetilde{\mathcal{H}} \to E(a+x) \otimes \widetilde{\mathcal{H}} \equiv \widetilde{\mathcal{H}}$. This proves our claim.

As the set $\{\theta_a(S)\xi : S \in E(a), \xi \in \widetilde{\mathcal{H}}\}\$ is total in $\widetilde{\mathcal{H}}$ and $\{\theta_x\}_{x \in \Omega}$ is anti multiplicative, it follows that $\widetilde{\theta}_x = \theta_x$ if $x \in \Omega$. Let $x \in P, y \in \Omega, T \in E(x)$ and $S \in E(y)$ be given. For $R \in E(a)$ and $\xi \in \widetilde{\mathcal{H}}$, calculate as follows to observe that

$$\begin{split} \widetilde{\theta}_{x+y}(TS)\theta_a(R)\xi &= \theta_{(a+x)+y}(RTS)\xi \\ &= \theta_y(S)\theta_{a+x}(RT)\xi \\ &= \theta_y(S)\widetilde{\theta_x}(T)\theta_a(R)\xi. \end{split}$$

Hence it follows that

$$\widetilde{\theta}_{x+y}(TS) = \theta_y(S)\widetilde{\theta}_x(T)$$
 (2.2)

for $x \in P$, $y \in \Omega$, $T \in E(x)$ and $S \in E(y)$.

Let $x, y \in P$, $T \in E(x)$ and $S \in E(y)$ be given. Let $R \in E(a)$ be of unit norm. Calculate as follows to observe that

$$\begin{aligned} \theta_a(R)\widetilde{\theta}_{x+y}(TS) &= \theta_{x+y+a}(TSR) \text{ (by Eq. 2.2)} \\ &= \theta_{x+(y+a)}(T(SR)) \\ &= \theta_{y+a}(SR)\widetilde{\theta}_x(T) \text{ (by Eq. 2.2)} \\ &= \theta_a(R)\widetilde{\theta}_y(S)\widetilde{\theta}_x(T) \text{ (by Eq. 2.2)}. \end{aligned}$$

Premultiplying the above equation by $\theta_a(R)^*$, we get

$$\widetilde{\theta}_{x+y}(TS) = \widetilde{\theta}_y(S)\widetilde{\theta}_x(T) \tag{2.3}$$

for $x, y \in P$, $T \in E(x)$ and $S \in E(y)$.

Note that for $x \in P$, the map $E(x) \ni T \to \widetilde{\theta}_x(T) \in B(\widetilde{\mathcal{H}})$ is linear and norm preserving. A direct calculation reveals that for $x \in P$, $T_1, T_2 \in E(x)$,

$$\widetilde{\theta}_x(T_2)^*\widetilde{\theta}_x(T_1) = \langle T_1|T_2\rangle_{E(x)}.$$
(2.4)

For $x \in P$, let $\widetilde{F}(x) := \{\widetilde{\theta}_x(T) : T \in E(x)\}$. Fix $x \in P$. It follows from Eq. 2.4 that there exists a unique normal *-endomorphism denoted $\widetilde{\beta}_x$ on $B(\widetilde{\mathcal{H}})$ such that the intertwining space of $\widetilde{\beta}_x$ is $\widetilde{F}(x)$. Eq. 2.3 implies that the family $\widetilde{\beta} := \{\widetilde{\beta}_x\}_{x \in P}$ forms a semigroup of endomorphisms.

Note that $\widetilde{F}(x) = F(x)$ for every $x \in \Omega$. Hence $\widetilde{\beta}_x = \beta_x$ for $x \in \Omega$. The semigroup $\widetilde{\beta}$ agrees with an E_0 -semigroup β on the interior Ω . Thanks to Lemma 4.1 of [6], it follows that $\widetilde{\beta}$ is an E_0 -semigroup. Since Ω is dense in P, it follows that $\widetilde{\beta}_x = \beta_x$ for every $x \in P$. Consequently, we have $\widetilde{F}(x) = F(x)$ for every $x \in P$. This completes the proof.

Proof of Proposition 2.1: Let $\{\widetilde{\theta}_x\}_{x\in P}$ be a family of unitary operators as in Lemma 2.3. For $x\in P$, denote the set of decomposable vectors in E(x) by D(x). A moment's reflection on the definition shows that the decomposable vectors of F(x) is $\{\widetilde{\theta}_x(T): T\in D(x)\}$. The conclusion is now immediate.

3 A counterexample

In this section, we produce the promised counterexample, i.e. a CCR flow which is asymmetric. Let us recall the definition of a CCR flow associated to a pure isometric representation. Suppose $V: P \to B(\mathcal{H})$ is a pure isometric representation. Recall that V is said to be pure if $\bigcap_{x \in P} V_x \mathcal{H} = \{0\}$. Denote the symmetric Fock space of \mathcal{H} by $\Gamma(\mathcal{H})$.

The CCR flow associated to V, denoted $\alpha^V := \{\alpha_x\}_{x \in P}$, is the unique E_0 -semigroup on $B(\Gamma(\mathcal{H}))$ such that the following equation is satisfied. For $x \in P$ and $\xi \in \mathcal{H}$,

$$\alpha_x(W(\xi)) = W(V_x \xi)$$

where $\{W(\xi): \xi \in \mathcal{H}\}$ is the set of Weyl operators on $\Gamma(\mathcal{H})$. For more details regarding multiparameter CCR flows, we refer the reader to [1] and [2].

Remark 3.1 A few remarks are in order.

- (1) In [7], a strongly continuous isometric representation, indexed by Ω , is constructed out of a decomposable E_0 -semigroup (see Proposition 4.1 of [7]). Moreover the resulting isometric representation, up to unitary equivalence, is a cocycle conjugacy invariant.
- (2) It is shown in [7] that the CCR flow α^V is decomposable. Moreover the isometric representation constructed out of the decomposable E_0 -semigroup α^V is V itself (see Proposition 5.1 of [7]).

Fix a pure isometric representation V of P on a Hilbert space \mathcal{H} . Denote the CCR flow α^V by α and its opposite by α^{op} . Denote the isometric representation constructed out of α^{op} by V^{op} . If we unwrap all the details regarding the construction of V^{op} , which is routine, we see that V^{op} has the following explicit description. Denote the Hilbert space on which V^{op} acts by \mathcal{H}^{op} .

Define an equivalence relation on $\{(\xi, a) : \xi \in Ker(V_a^*), a \in \Omega\}$ as follows. We say $(\xi, a) \sim (\eta, b)$ if and only if $V_b \xi = V_a \eta$. Let H^{op} be the set of equivalence classes. Then H^{op} has an inner product space structure where the addition, scalar multiplication and inner product are given by

$$[(\xi, a)] + [(\eta, b)] = [(V_b \xi + V_a \eta, a + b)]$$
$$\lambda[(\xi, a)] = [(\lambda \xi, a)]$$
$$\langle [(\xi, a)] | [(\eta, b)] \rangle = \langle V_b \xi | V_a \eta \rangle.$$

Then \mathcal{H}^{op} is the completion of H^{op} and for $a \in \Omega$, the operator V_a^{op} is given by the equation

$$V_a^{op}[(\xi, b)] = [(\xi, a + b)].$$

To produce a counter example of a CCR flow which is asymmetric, it suffices to construct a pure isometric representation V such that V and V^{op} are not unitarily equivalent. For if $\alpha := \alpha^V$ and α^{op} are cocycle conjugate then by Remark 3.1, it follows that V and V^{op} are unitarily equivalent.

First we obtain a better description of V^{op} . Let U be the minimal unitary dilation of V. More precisely, there exists a Hilbert space $\widetilde{\mathcal{H}}$ containing \mathcal{H} as a closed subspace and a strongly continuous unitary representation $U := \{U_x\}_{x \in \mathbb{R}^d}$ on $\widetilde{\mathcal{H}}$ such that the following conditions are satisfied.

- (1) For $a \in \Omega$ and $\xi \in \mathcal{H}$, $U_a \xi = V_a \xi$, and
- (2) the increasing union $\bigcup_{a\in\Omega} U_a^*\mathcal{H}$ is dense in $\widetilde{\mathcal{H}}$.

The minimal unitary dilation is unique up to unitary equivalence and the existence of such a dilation is given by an inductive limit procedure.

Set $\mathcal{K} := \mathcal{H}^{\perp}$. Since \mathcal{H} is invariant under $\{U_a\}_{a \in \Omega}$, it follows that \mathcal{K} is invariant under $\{U_{-a}: a \in \Omega\} = \{U_a^*: a \in \Omega\}.$ For $a \in \Omega$, let W_a be the operator on \mathcal{K} which is the restriction of U_{-a} . Then $W := \{W_a\}_{a \in \Omega}$ is a strongly continuous semigroup of isometries on \mathcal{K} .

Proposition 3.2 With the foregoing notation, we have the following.

- (1) The isometric representation W is pure, i.e. $\bigcap_{a\in\Omega}W_a\mathcal{K}=\{0\}.$
- (2) The isometric representations W and V^{op} are unitarily equivalent.

Proof. Fix a point $a \in \Omega$. Set $S := V_a$. Recall the following Archimedean property. Given $x \in \mathbb{R}^d$, there exists a positive integer n such that na > x (see Lemma 3.1 of [5]).

Thus $\{0\} = \bigcap_{b \in \Omega} V_b \mathcal{H} = \bigcap_{n \geq 1} V_{na} \mathcal{H} = \bigcap_{n \geq 1} S^n \mathcal{H}$. In other words, S is a pure isometry. By the Archimedean prinicple, we have the equaltiy $\bigcup_{b \in \Omega} U_b^* \mathcal{H} = \bigcup_{n \geq 1} U_a^{*n} \mathcal{H}$. Hence $\bigcup U_a^{*n}\mathcal{H}$ is dense in $\widetilde{\mathcal{H}}$. In otherwords, the discrete one parameter group of unitaries

 $\{\overline{U}_{na}:n\in\mathbb{Z}\}$ is the minimal unitary dilation of the discrete one parameter isometric representation $\{S^n\}_{n>0}$.

Using Wold decomposition, we can identify \mathcal{H} with $\ell^2(\mathbb{N}) \otimes K$ for some Hilbert space K and S with the standard one sided shift with multiplicity. Then $\widetilde{\mathcal{H}}$ can be identified with $\ell^2(\mathbb{Z}) \otimes K$ with U_a identified with the bilateral shift with multiplicity. Once this

identification is made, it is clear that $\bigcap_{n\geq 0} W_{na}\mathcal{K} = \bigcap_{n\geq 0} U_a^{*n}\mathcal{K} = \{0\}$. Once again by the Archimedean principle, we have the equality $\bigcap_{b\in\Omega} W_b\mathcal{K} = \bigcap_{n\geq 0} W_{na}\mathcal{K} = \{0\}$. This proves (1).

Let $a \in \Omega$. We claim that the image of the map

$$Ker(W_a^*) \ni \xi \to U_a \xi \in \widetilde{\mathcal{H}}$$

is contained in $Ker(V_a^*)$. Let $\xi \in Ker(W_a^*)$ and $\eta \in \mathcal{K}$ be given. Observe that

$$\langle U_a \xi | \eta \rangle = \langle \xi | U_a^* \eta \rangle = \langle \xi | W_a \eta \rangle = \langle W_a^* \xi | \eta \rangle = 0.$$

This proves that for $\xi \in Ker(W_a^*)$, $U_a\xi \in \mathcal{H}$. Let $\xi \in Ker(W_a^*)$ and $\eta \in \mathcal{H}$ be given. Using the fact that $\xi \in \mathcal{K}$ and $\eta \in \mathcal{H}$, observe that

$$\langle U_a \xi | V_a \eta \rangle = \langle U_a \xi | U_a \eta \rangle = \langle \xi | \eta \rangle = 0.$$

Thus $U_a\xi$ is orthogonal to $Ran(V_a)$. Coupled with the fact that $U_a\xi \in \mathcal{H}$, we conclude that $U_a\xi \in Ker(V_a^*)$. This proves our claim. A calculation similar to the one above implies that the image of the map $Ker(V_a^*) \ni \eta \to U_{-a}\eta \in \widetilde{\mathcal{H}}$ is contained in $Ker(W_a^*)$. Consequently the map $Ker(W_a^*) \ni \xi \to U_a\xi \in Ker(V_a^*)$ is a unitary.

By (1), we have $\mathcal{K} = \overline{\bigcup_{a \in \Omega} Ker(W_a^*)}$. Note that the family of inner product preserving maps

$$\left\{Ker(W_a^*)\ni\xi\to[(U_a\xi,a)]\in\mathcal{H}^{op}\right\}_{a\in\Omega}$$

patch together and defines a unitary map from $\mathcal{K} = \overline{\bigcup_{a \in \Omega} Ker(W_a^*)}$ to \mathcal{H}^{op} , which we denote by T, such that the following holds. If $\xi \in Ker(W_a^*)$ for some $a \in \Omega$ then

$$T\xi = [(U_a\xi, a)].$$

It is clear that T intertwines the isometric representations W and V^{op} . This proves (2). The proof is now complete.

Let $A \subset \mathbb{R}^d$ be such that A is non-empty, proper, closed and $A + P \subset A$. Such subsets were called P-modules in [2]. Consider the Hilbert space $L^2(A)$. For $x \in P$, let V_x be the isometry on $L^2(A)$ defined by the equation

$$V_x(f)(y) := \begin{cases} f(y-x) & \text{if } y-x \in A, \\ 0 & \text{if } y-x \notin A. \end{cases}$$

$$(3.5)$$

Then $(V_x)_{x\in P}$ is an isometric representation of P which we denote by V^A . We call V^A the isometric representation associated to the P-module A. Moreover the representation V^A is pure. In what follows, Int(A) denotes the interior of A.

Lemma 3.3 Let A be a P-module. We have the following.

- (1) The increasing union $\bigcup_{a \in \Omega} (Int(A) a) = \mathbb{R}^d$.
- (2) Given a compact subset $K \subset \mathbb{R}^d$, there exists $a \in \Omega$ such that K is contained in Int(A) a.
- (3) The minimal unitary dilation of V^A is the left regular representation of \mathbb{R}^d on $L^2(\mathbb{R}^d)$.

Proof. Since A is a P-module, it is clear that if a < b with $a, b \in \Omega$ then Int(A) - a is contained in Int(A) - b. By translating, if necessary, we can assume $0 \in A$. Hence $P \subset A$ and $\Omega \subset Int(A)$. Observe the equality $\mathbb{R}^d = \Omega - \Omega = \bigcup_{a \in \Omega} (\Omega - a) \subset \bigcup_{a \in \Omega} (Int(A) - a)$. This proves (1).

Fix an interior point $a \in \Omega$. By (1) and by the Archimedean property, it follows that $(Int(A) - na)_{n\geq 1}$ is an increasing sequence of open sets which increases to \mathbb{R}^d . Now (2) is immediate.

Let $\{U_x\}_{x\in\mathbb{R}^d}$ be the left regular representation of \mathbb{R}^d on $L^2(\mathbb{R}^d)$. From the definition it follows that for $a\in\Omega$, V_a is the compression of U_a onto $L^2(A)$. Observe that for $a\in\Omega$, $U_a^*L^2(A)=L^2(A-a)$. It follows from (2) that $\bigcup_{a\in\Omega}U_a^*L^2(A)$ contains the space

of continuous functions with compact support. Consequently $\bigcup_{a\in\Omega}U_a^*L^2(A)$ is dense in

 $L^2(\mathbb{R}^d)$. Hence $\{U_x\}_{x\in\mathbb{R}^d}$ has all the properties required to be the minimal unitary dilation of V^A . This completes the proof.

Note that if A is a P-module then $-(IntA)^c$ is a P-module. Fix a P-module A and let $V := V^A$ be the isometric representation associated to A. Set $B := -(Int(A))^c$.

Proposition 3.4 Keep the foregoing notation.

- (1) The isometric representation V^{op} is unitarily equivalent to the representation V^{B} .
- (2) The representations V and V^{op} are unitarily equivalent if and only if there exists $z \in \mathbb{R}^d$ such that A = B + z.

Proof. Thanks to Lemma II.12 of [4], A and Int(A) differ by a set of measure zero. Thus we can identify $L^2(A)$ with $L^2(Int(A))$. Lemma 3.3 and Proposition 3.2 implies that V^{op} is equivalent to the isometric representation $W = \{W_a\}_{a \in \Omega}$ acting on the Hilbert

space $L^2((Int(A))^c)$ where the operators $\{W_a\}_{a\in\Omega}$ are given by the following equation. For $a\in\Omega$ and $f\in L^2((Int(A))^c)$

$$W_a(f)(y) := \begin{cases} f(y+a) & \text{if } y+a \in (Int(A))^c, \\ 0 & \text{if } y+a \notin (Int(A))^c. \end{cases}$$

$$(3.6)$$

The inversion $\mathbb{R}^d \ni x \to -x \in \mathbb{R}^d$ induces a unitary between the Hilbert spaces $L^2((Int(A))^c)$ and $L^2(B)$ and intertwines the representations W and V^B . This proves (1).

It is clear that if A is a translate of B then $V = V^A$ and $V^{op} = V^B$ are unitarily equivalent. On the other hand, suppose V^A and V^B are unitarily equivalent. Then the associated CCR flows α^{V^A} and α^{V^B} are cocycle conjugate. By Theorem 1.2 of [2], it follows that A and B are translates of each other. This completes the proof.

Thus to produce a counterexample of a CCR flow which is not cocycle conjugate to its opposite, it suffices to produce a P-module A such that A is not a translate of $-(Int(A))^c$. The cone P itself is one such candidate. Recall that we have assumed $d \geq 2$. Let us recall the notion of extreme points. For a subset C of \mathbb{R}^d and a point $x \in C$, we say x is an extreme point of C if $x = \frac{y+z}{2}$ with $y, z \in C$ then y = z = x.

Lemma 3.5 The sets P and $-\Omega^c$ are not translates of each other.

Proof. First we claim that $P \cup -P \neq \mathbb{R}^d$. Suppose not. Since we have assumed that $P \cap -P = \{0\}$, it follows that the only boundary point of P is $\{0\}$. Fix $a \in \Omega$. Proposition 2.3 of [2] implies that the map

$$\partial(P)\times(0,\infty)\ni(x,s)\to x+sa\in\Omega$$

is a homeomorphism. But $\partial(P) = \{0\}$ and hence $\Omega = \{sa : s > 0\}$. Since Ω spans \mathbb{R}^d , it follows that d = 1 contradicting our assumption. Therefore $P \cup -P \neq \mathbb{R}^d$.

Note that the set of extreme points of P is $\{0\}$. For we have assumed that P contains no line. On the other hand, we claim that Ω^c has no extreme point. Note that $t\Omega^c \subset \Omega^c$ for t > 0. Hence the set of extreme points of Ω^c is contained in $\{0\}$. But 0 is not an extreme point of Ω^c . To see this, pick $x \notin P \cup -P$. Then $x \in \Omega^c$, $-x \in \Omega^c$ and $x \neq 0$. But $0 = \frac{x + (-x)}{2}$. This proves that the set of extreme points of Ω^c is empty. Consequently, the set of extreme points of any translate of $-\Omega^c$ is empty. Hence P and $-\Omega^c$ are not translates of each other. This completes the proof.

¹See also Page 26 of [7].

Remark 3.6 Let V be a pure isometric representation of P and $\alpha = \alpha^V$ be the associated CCR flow. Let V^{op} be the isometric representation corresponding to the decomposable E_0 -semigroup α^{op} . Since α is spatial, it follows that α^{op} is spatial. (Recall that an E_0 -semigroup is said to be spatial if its product system admits a nowhere vanishing multiplicative measurable cross section). By Theorem 4.4 of [7], it follows that α^{op} and $\alpha^{V^{op}}$ are cocycle conjugate. Thus, to summarise, an opposite of a CCR flow is a CCR flow but not necessarily the same as the original one.

References

- [1] Arjunan Anbu, R. Srinivasan, and S. Sundar, *E-semigroups over closed convex cones*, arxiv/math.OA:1807.11375.
- [2] Arjunan Anbu and S. Sundar, *CCR flows associated to closed convex cones*, to appear in Muenster Journal of Mathematics, arxiv/math.OA:1901.00265.
- [3] William Arveson, *Noncommutative dynamics and E-semigroups*, Springer Monographs in Mathematics, Springer-Verlag, New York, 2003.
- [4] Joachim Hilgert and Karl-Hermann Neeb, Wiener-Hopf operators on ordered homogeneous spaces. I, J. Funct. Anal. 132 (1995), no. 1, 86–118.
- [5] S. P. Murugan and S. Sundar, On the existence of E_0 -semigroups—the multiparameter case, Infin. Dimens. Anal. Quantum Probab. Relat. Top. **21** (2018), no. 2, 1850007, 20.
- [6] S.P. Murugan and S. Sundar, E_0^P -semigroups and Product systems, arxiv/math.OA:1706.03928.
- [7] S. Sundar, Arveson's characterisation of CCR flows: the multiparameter case, arxiv/math.OA:1906:05493v2.
- [8] Boris. Tsirelson, From random sets to continuous tensor products: answers to three questions of W. Arveson, arxiv/math.FA:0001070.
- [9] Boris Tsirelson, Non-isomorphic product systems, Advances in quantum dynamics (South Hadley, MA, 2002), Contemp. Math., vol. 335, Amer. Math. Soc., Providence, RI, 2003, pp. 273–328.

$S. \ Sundarsobers {\tt @gmail.com})$

Institute of Mathematical Sciences (HBNI), CIT Campus,

Taramani, Chennai, 600113, Tamilnadu, INDIA.