# The Role of High-mass Stellar Binaries in the Formation of High-mass Black Holes in Dense Star Clusters

Ambreesh Khurana  $\textcircled{D}^1$  and Sourav Chatterjee  $\textcircled{D}^1$ 

<sup>1</sup>Department of Astronomy & Astrophysics, Tata Institute of Fundamental Research, Homi Bhabha Road, Navy Nagar, Colaba, Mumbai 400005, India

# ABSTRACT

Recent detections of gravitational waves from mergers of binary black holes (BBHs) with pre-merger source-frame individual masses in the so-called upper mass-gap, expected due to (pulsational) pair instability supernova ((P)PISN), have created immense interest in the astrophysical production of highmass black holes (BHs). Previous studies show that high-mass BHs may be produced via repeated BBH mergers inside dense star clusters. Alternatively, inside dense star clusters, stars with unusually low core-to-envelope mass ratios can form via mergers of high-mass stars, which then can avoid (P)PISN, but produce high-mass BHs via mass fallback. We simulate detailed star-by-star multi-physics models of dense star clusters using the Monte Carlo cluster evolution code, CMC, to investigate the role of primordial binary fraction among high-mass stars ( $\geq 15 M_{\odot}$ ) on the formation of high-mass BHs. We vary the high-mass stellar binary fraction  $(f'_{h,15})$  while keeping all other initial properties, including the population of high-mass stars, unchanged. We find that the number of high-mass BHs, as well as the mass of the most massive BH formed via stellar core-collapse are proportional to  $f'_{b,15}$ . In contrast, there is no correlation between  $f'_{b,15}$  and the number of high-mass BHs formed via BH-BH mergers. Since the total production of high-mass BHs is dominated by BH-BH mergers in old clusters, the overall number of high-mass BHs produced over the typical lifetime of globular clusters is insensitive to  $f'_{b,15}$ . Furthermore, we study the differences in the demographics of BH-BH mergers as a function of  $f'_{b,15}$ .

### 1. INTRODUCTION

Isolated high-mass stars with main-sequence (MS) mass  $\gtrsim 100 M_{\odot}$  (He core mass  $\gtrsim 40 M_{\odot}$ ), during their final stages of evolution, can reach core temperature and pressure high enough to initiate electron-positron pair production (e.g., Rakavy & Shaviv 1967; Heger & Woosley 2002). This leads to a sudden decrease in pressure and the core collapses. The resulting thermonuclear explosion(s) due to oxygen burning can either cause the star to lose mass in episodes, resulting in the so-called pulsational pair instability supernova (PPISN) before collapsing to a black hole (BH) whose mass is thus limited, or destroy the star completely, leaving behind no remnant at all as a result of the so-called pair instability supernova (PISN; e.g., Barkat et al. 1967; Woosley et al. 2002). One the other hand, models suggest, very high-

souravchatterjee.tifr@gmail.com

mass stars with an initial mass  $\gtrsim 260 M_{\odot}$  (He core mass  $\gtrsim 130 M_{\odot}$ ) can again collapse to create BHs with mass  $M_{\rm BH} \gtrsim 100 M_{\odot}$  due to photo-disintegration in the core (e.g., Heger et al. 2003; Woosley et al. 2002). These processes are thus expected to create a gap between ~ 40 and ~ 100  $M_{\odot}$  in the birth mass function of BHs (e.g., Heger & Woosley 2002), although, the exact boundaries of this so-called "upper mass-gap" are still under investigation (e.g., Farmer et al. 2019, 2020; Costa et al. 2021; Vink et al. 2021; Farag et al. 2022).

Interestingly, gravitational-wave (GW) observatories have detected binary BH (BBH) merger events, such as GW190521 (Abbott et al. 2020), GW190403\_051519, GW190426\_190642 (Abbott et al. 2024), GW200220\_061928 (Abbott et al. 2023), and GW200129\_114245 (Nitz et al. 2023), with individual pre-merger source-frame masses well within the upper mass-gap. These detections have created intense interest in exploring the physical processes that could lead to the formation of such high-mass BHs. Inside dense star clusters, BH-BH binaries are assembled via dynam-

ambreesh.sci@gmail.com

ical processes which then can merge via the emission of GWs inside the cluster or after getting ejected from the cluster (e.g., Banerjee et al. 2010; Rodriguez et al. 2015, 2016; Askar et al. 2017; Banerjee 2017; Chatterjee et al. 2017a,b; Banerjee 2018a,b; Samsing et al. 2018; Kremer et al. 2020). If the merger product is not ejected from the cluster, the newly formed BH from the BBH merger may acquire another BH companion via the same dynamical processes and merge again (e.g., Rodriguez et al. 2018; Rodriguez et al. 2020; Martinez et al. 2020). This process can continue until the merger product is ejected from the cluster either via recoil from dynamical interactions or merger-driven kicks. It has been proposed that BH-BH mergers involving BHs within the upper mass gap may have been formed via these socalled higher generation mergers (e.g., Rodriguez et al. 2019; Weatherford et al. 2021).

On the other hand, inside dense star clusters, at early times, high-mass stars may collide with other stars. These collision products can have a high enough mass to form a mass-gap BH but have unusually low core-toenvelope mass ratios (e.g., Ballone et al. 2022) such that the core mass is not high enough to initiate (P)PISN. Thus, the star does not lose too much mass during corecollapse. This could create a BH more massive than what is possible via single star evolution as the initial BH could grow in mass via at least partial fallback of the unusually massive envelope (e.g., Di Carlo et al. 2019; Spera et al. 2019; Di Carlo et al. 2020; Kremer et al. 2020; Renzo et al. 2020; González et al. 2021; Costa et al. 2022). Note that both these processes, i.e., BH-BH mergers and core-collapse of stellar merger products, may undergo within dense star clusters and the two processes are not necessarily independent (e.g., Sedda et al. 2023; Prieto et al. 2022). For example, a high-mass BH, formed via the latter mechanism, may later take part in multiple BH-BH mergers via the former pathway. It is interesting, however, to understand the relative importance of these processes as a function of the initial cluster properties and age.

Theoretical models and observations suggest that high-mass stars are often in binaries or higher-order multiples (e.g., Sana et al. 2012; Offner et al. 2023). Recently, González et al. (2021) (hereafter GKC21) showed that massive star clusters, modeled using the Cluster Monte Carlo (CMC) code (Joshi et al. 2000, 2001; Fregeau & Rasio 2007; Chatterjee et al. 2010; Goswami et al. 2012; Morscher et al. 2013; Pattabiraman et al. 2013; Breivik et al. 2020; Rodriguez et al. 2022), with an initial binary fraction among high-mass stars (> 15  $M_{\odot}$ ),  $f_{b,15} = 1$ , can produce BHs with mass in the upper mass-gap and even higher (collectively, "high-

mass BHs", hereafter) via core-collapse of stellar merger products. In contrast, models with  $f_{b,15} = 0$ , produced almost no high-mass BHs (GKC21, in particular, see their Figure 2).<sup>1</sup> There were two important differences between the two sets of models with  $f_{b,15} = 0$  and 1. First, all high-mass stars in  $f_{b,15} = 1(0)$  models were in binaries (singles). The other, relatively less apparent, difference is in the total initial number of high-mass stars. In CMC, the masses of stars, primaries in case of binaries  $(M_{\text{prim}})$ , are sampled assuming an initial stellar mass function (IMF). Then, depending on the assumed overall and high-mass binary fractions,  $f_{\rm b}$  and  $f_{{\rm b},15}$ , a randomly chosen fraction of these stars (or primaries) are assigned a binary companion with mass  $(M_{sec})$  based on the adopted mass-ratio  $(q \equiv M_{\rm sec}/M_{\rm prim})$  distribution and range. Thus, for a fraction of the companions,  $M_{\rm sec} > 15 M_{\odot}$ . As a result, models with  $f_{\rm b.15} = 1$ end up with a higher number of high-mass stars than the models with  $f_{b,15} = 0$ . In other words, changing  $f_{b,15}$  does not conserve the overall IMF. For example, using the adopted initial conditions in GKC21, i.e.,  $N = 8 \times 10^5$  (number of initial objects, singles and binaries), stellar IMF based on Kroupa (2001), uniform mass ratio distribution for high-mass stars between  $q_{15} = 0.6$  and 1, and  $f_{b,15} = 1$ , the total number of initial stars with mass  $\geq 15 M_{\odot}$ ,  $N_{15} = 3782^{+130}_{-63}$ . In contrast, using the same assumptions, but with  $f_{b,15} = 0$ ,  $N_{15} = 2181^{+74}_{-32}$ . This leads to a difference in initial  $N_{15}$  of  $1606^{+53}_{-44}$ . Interestingly, GKC21 also found that most ( $\approx 96\%$ ) high-mass BHs formed via stellar collisions during dynamical encounters as opposed to binary stellar evolution. Thus, it is not clear whether the formation of much higher numbers of high-mass BHs in their  $f_{b,15} = 1$  models is because of the difference in  $f_{\rm b,15}$  or  $N_{15}$ . Moreover, given the importance of highmass stars in the overall evolution and even survival of the host cluster and the formation of stellar exotica (e.g., Chatterjee et al. 2017b) disentangling the effects of the differences in  $N_{15}$  and  $f_{b,15}$  is very interesting.

In this paper, we investigate whether the high-mass BHs that form in star cluster models with  $f_{b,15} = 1$  indeed require those high-mass stars to be in a binary initially or this is simply a result of a higher  $N_{15}$  in these models compared to those with  $f_{b,15} = 0$ . We create models systematically varying the initial fraction of high-mass stars in binaries between 0 and 1 keeping  $N_{15}$ 

<sup>&</sup>lt;sup>1</sup>  $f_{b,15} \equiv N_{b,15}/(N_{b,15} + N_{s,15})$ , where  $N_{b,15}$  and  $N_{s,15}$  represent the number of binaries containing at least one high-mass star and the number of high-mass single stars, respectively. Similarly,  $f_b \equiv N_b/(N_b + N_s)$ , where  $N_b$  ( $N_s$ ) denote the number of lowmass binaries (single).

and other initial properties fixed. We investigate the long-term evolution of these model clusters paying particular attention to the formation of high-mass BHs via core-collapse of stellar merger products as well as BH-BH mergers. Furthermore, we investigate the relative contribution to high-mass BH-BH mergers from these two channels as a function of the cluster age.

In section 2, we describe our star cluster models. We also describe the process by which we initialise cluster models with varying  $f_{b,15}$  keeping  $N_{15}$  and other properties unchanged. In section 3, we present our findings in detail. Finally, we conclude and provide a summary in section 4.

### 2. METHODS

We use the highly realistic Hénon-type Monte Carlo cluster evolution code, CMC (Rodriguez et al. 2022) to simulate star clusters. The stellar evolution in CMC is handled by the population synthesis code, COSMIC (Breivik et al. 2020). As the so-called 'control group' of our cluster models, we choose the initial parameters as follows. This group of models has initial  $f_{b,15} = 1$  and the binary fraction in lower-mass stars,  $f_{\rm b} = 0.05$ . The initial  $N = 8 \times 10^5$ . The single and primary stars are sampled from a Kroupa (2001) IMF ranging from 0.08 to  $150 M_{\odot}$ . For the binaries, eccentricities (e) are assumed to be thermal (dN/de = 2e, Jeans 1919) while the orbital periods (P) are drawn from the distribution,  $dN/d\log P \propto P^{-0.55}$  (Sana et al. 2012) between  $\log (P/day) = 0.15$  and the smaller of the hard-soft boundary and  $\log (P/day) = 5.5$ . The mass ratio between binary members is sampled from a uniform distribution between q = 0.1 and 1 for lower-mass primaries and  $q_{15} = 0.6$  and 1 for the high-mass primaries. This results in a total initial cluster mass  $M_{\rm cl}/M_{\odot} \sim 5.4 \times 10^5$ . The cluster members are distributed according to the King (1966) profile with a concentration parameter,  $w_0 = 5$  and a virial radius,  $r_v = 1 \text{ pc.}$  The metallicity of the stars is fixed at  $0.1 Z_{\odot}$ , where  $Z_{\odot} = 0.02$ . We simulate each cluster up to  $\approx 13.8 \,\text{Gyr}$ . In order to take statistical fluctuations into account, we run 10 models with the same properties but generated using different random seeds.

To explore the importance of initial  $f_{b,15}$  compared to  $N_{15}$  in creating high-mass BHs, we create additional cluster models which have the same initial  $N_{15}$  as the control group but different  $f_{b,15}$ . For each model in our control group with  $f_{b,15} = 1$ , we randomly choose 25%, 50%, 75%, and *all* high-mass stellar binaries and break them into singles. We redistribute the new collection of single and binary stars in the cluster according to the same King (1966) profile. This leaves us with a total number of stellar objects (binary/single) higher than  $8 \times 10^5$ . This procedure helps us create sets of initial cluster models with  $f'_{\rm b,15}$  = 0.75, 0.5, 0.25, and 0, respectively, where the individual stellar populations remain identical between corresponding cluster models. Here,  $f'_{b,15}$  denotes the fraction of initially high-mass binaries that we leave intact. The changed notation indicates that  $f'_{b,15}$  and  $f_{b,15}$  are not equal unless  $f'_{b,15} = 0$ or 1 (Table 1). If the production of high-mass BHs requires enough number of high-mass stars in the cluster independent of whether they are initially in a binary or not, then these cluster models with different initial  $f'_{b,15}$ but the same  $N_{15}$  would show little statistical difference. On the other hand, if  $f'_{b,15}$  is indeed a critical property, then we would find systematic differences between these groups of models.

The formation of high-mass BHs via core-collapse of stellar merger products depends on how stellar collisions are treated in the models. GKC21 used the simple 'sticky sphere' approximation for MS and giant star collisions (section 2, Kremer et al. 2020). In contrast, we use the updated prescriptions for collisions involving giant stars currently used in CMC (Ye et al. 2022). These updates are motivated by the expectation that collisions involving giant stars should be better represented by something similar to a common envelope (CE) evolution in which the core of the giant and the companion (a MS star or a compact object) spiral towards each other losing orbital energy to the giant's envelope that enshrouds them. If the envelope can be ejected due to the energy released by the shrinking orbit before they merge, a binary should form composed of the giant's core and the secondary object. If both of the colliding stars are giants, the binary formed would be composed of the cores of the two giants. However, if the envelope can not be ejected, they merge. The updated prescription calculates orbital parameters for the final binary assuming a CE evolution (Hurley et al. 2002) as well as from a set of parameterised equations (Ivanova et al. 2006) adapted from the results of smoothed particle hydrodynamics simulations (Lombardi, Jr. et al. 2006). If not merged, the final binary is assigned the orbital properties corresponding to the smaller semi-major axis (a) from the above two solutions. If the formed binary satisfies the condition for Roche-lobe overflow, the participating stars are merged including the envelope (for more details on the updated prescription, see Ye et al. 2022).

In addition to the treatment of stellar collisions, our results also depend on the adopted prescriptions for modeling BH natal kicks and BH-BH merger recoil because these affect the retention of BHs in the cluster




Figure 1. Distribution of the average number of BHs produced via stellar core-collapse per model as a function of BH mass. Errorbars represent  $1\sigma$ . The vertical shaded region represents the upper mass-gap due to (P)PISN. Blue, orange, green, red, and purple denote models with  $f'_{b,15} = 1$ , 0.75, 0.5, 0.25, and 0, respectively. Although the  $M_{\rm BH}$  distributions in cluster models with different  $f'_{b,15}$  are similar below the upper mass-gap, the number of high-mass BHs produced depends on  $f'_{b,15}$ .

after they form and after a GW-driven merger, respectively. We assume that BHs receive natal kicks scaled down by the mass fallback fraction (Fryer et al. 2012) from the standard neutron star (NS) kicks, where NS natal kicks are drawn from a Maxwellian distribution with dispersion  $265 \text{ km s}^{-1}$  (Hobbs et al. 2005; Kremer et al. 2020). We also assume that BHs are born nonspinning (Breivik et al. 2020). The mass, spin, and GW recoil of the GW-driven merger products are estimated using the prescriptions given in Gerosa & Kesden (2016) based on numerical relativity calculations.<sup>2</sup>

In our simulations, we choose the prescription for (P)PISN physics parameterised from the simulations of Marchant et al. (2019), available by default in CMC-COSMIC (Breivik et al. 2020). This corresponds to an upper mass-gap in the range  $\approx 44.3 - 123 M_{\odot}$ . Furthermore, we use the "delayed" supernova (SN) explosion model for determining the mass of the remnant after SN (Fryer et al. 2012).

#### 3. RESULTS




Figure 2. Fraction of time the high-mass  $(M_{\star}/M_{\odot} > 15)$  stars exist as a part of a binary during the first 35 Myr, before either collapsing to a compact object, colliding with another star, or getting ejected from the cluster. Blue and orange denote example models with  $f'_{b,15} = 1$  and 0. Solid and dashed denote time fractions spent as a binary overall, and with the primordial companion, respectively.

# 3.1. Evolution in Early Times

To study high-mass BH formation in dense star clusters via core-collapse of stellar collision products, we first focus on the early (< 35 Myr) stages of evolution. This is the time when high-mass stars exist as luminous objects before turning into remnants such as BHs and NSs. Figure 1 shows the histograms of the mass of all BHs ( $M_{\rm BH}$ ) produced via core-collapse in our models with various initial  $f'_{\rm b,15}$ . We find that all core-collapse BHs form within a cluster age,  $t_{\rm cl}/{\rm Myr}$  < 35 in our models. The vertical shaded region represents the upper mass-gap corresponding to the assumptions in our simulations. Clearly, the high-mass end of the BH mass spectrum is sensitive to the initial  $f'_{\rm b,15}$ - models with higher initial  $f'_{\rm b,15}$  produce more high-mass BHs via core-collapse despite the same  $N_{15}$  in all models.

Being part of a binary initially does not ensure that the binary stays intact. Hence, we investigate the fractions of times high-mass stars remain part of a binary during this early  $(t_{\rm cl}/{\rm Myr} < 35)$  evolution before forming remnants, colliding with another star, or getting ejected from the cluster. Figure 2 shows the distribution for the fraction of time the high-mass stars remain in a binary system. We find that if initially in a binary, the high-mass stars remain part of a binary most of the time. Moreover, most of the time, the companion is the primordial one. In contrast, the high-mass stars in the  $f'_{\rm b,15} = 0$  models rarely capture a companion so early in the evolution of the cluster.

 $<sup>^2</sup>$  See Rodriguez et al. (2018) Appendix A for more details and a typo correction in the remnant-mass expression.

$f_{\mathrm{b},15}'$	$f_{ m b,15}$	$N_{ m BH,high}$			$M_{ m BH,max}/M_{\odot}$			
		Core-collapse	BH-BH Merger	Mass Transfer	Core-collapse	BH-BH Merger	Mass Transfer	
1	1	$4^{+1.75}_{-1}$	$21.5^{+1.5}_{-3.5}$	$20^{+4}_{-5}$	$119.5^{+20.7}_{-41.4}$	$112.3^{+11.8}_{-7.8}$	$50.7^{+1.5}_{-2.1}$	
0.75	0.63	$3^{+1.5}_{-0.75}$	$20.5^{+2}_{-1.25}$	$15^{+3}_{-2}$	$88^{+7.9}_{-30}$	$117.3^{+8.3}_{-19.3}$	$51.8^{+2.9}_{-3.3}$	
0.5	0.36	$1.5_{-0.5}^{+0.5}$	$21.5^{+3.75}_{-4.25}$	$10^{+2}_{-1}$	$54.4^{+10.3}_{-3.9}$	$112.8^{+2.5}_{-10.2}$	$51.8^{+1.4}_{-1.0}$	
0.25	0.16	$1^{+0}_{-1}$	$18^{+2}_{-1.75}$	$4^{+1}_{-1}$	$51.9^{+7.4}_{-8.3}$	$116.5^{+1.6}_{-18.2}$	$48.1_{-0.5}^{+0.3}$	
0	0	$0^{+0}_{-0}$	$20^{+3}_{-5}$	$0^{+0}_{-0}$	$43.5_{-0.1}^{+0.1}$	$113.9^{+1.6}_{-3.5}$	N/A	

**Table 1.** Demographics of the BH populations created in our star cluster models with varying  $f'_{b,15}$ . For each  $f'_{b,15}$  we show the corresponding  $f_{b,15}$ . We show the number of BHs with mass in the upper mass gap or above  $(N_{\text{BH,high}})$  and the maximum BH mass  $(M_{\text{BH,max}})$ . For each of the above, we separately show BHs created via stellar core-collapse, BH-BH mergers, and mass transfer in a binary. The numbers and errorbars denote the median and the quartiles.

The collision rate for stars is proportional to the stellar number density  $(n_{\star})$  and the square of the stellar radius  $(R_{\star})$ . Hence, if the mass segregation timescale  $(t_{seg})$  is shorter than the MS lifetime  $(t_{\rm MS})$ , then the high-mass stars can sink to the high-density core as large objects significantly boosting the rate of physical collisions. In contrast, if  $t_{\rm seg} > t_{\rm MS}$ , then the high-mass stellar collisions are suppressed significantly. By the time high-mass stars can sink to the center, they evolve and form remnants. Formation of high-mass BHs via core-collapse of stellar collision products of course requires that the collisions happen before remnant formation (e.g., Di Carlo et al. 2019). The mass loss during remnant formation also leads to rapid overall expansion of the cluster leading to a sharp decrease in  $n_{\star}$  at the core (e.g., Chatterjee et al. 2017b). If a high-mass star is part of a binary, then  $t_{\text{seg}}$  reduces by a factor of  $\sim (1 + q_{15})$  thus increasing the chance that the high-mass star sinks to the core before remnant formation. Moreover, for a binary, the strong interaction cross-section is  $\propto a^2$ , significantly higher than single-star collision cross section. For the typical velocity dispersion inside dense stellar clusters, the dominant resonant type interactions often lead to physical collisions involving the large high-mass stars (e.g., Fregeau et al. 2004).

Indeed, we find a clear correlation between the number of high-mass BHs produced via core collapse and  $f'_{b,15}$ . While the models with  $f'_{b,15} = 1$  create  $\approx 4$  high-mass BHs via stellar core-collapse per cluster, this number decreases almost linearly to  $\approx 0$  for the models with  $f'_{b,15} < 0.1$  (Figure 3, middle). Clusters with higher  $f'_{b,15}$  are expected to bring in more high-mass stars to the core before remnant formation, which leads to a higher chance of high-mass BH production via corecollapse of collision products. We find that the mass of the most massive BH ( $M_{\rm BH,max}$ ) a cluster produces via core-collapse depends on  $f'_{b,15}$  as well (Figure 3, bottom). While models with  $f'_{b,15} = 1$  could produce BHs with mass as high as  $\approx 180 M_{\odot}$ , the heaviest BH in the  $f'_{b,15} = 0$  models is  $\approx 60 M_{\odot}$ . The median  $M_{\rm BH,max}$ 

decreases from  $\approx 120 M_{\odot}$  for  $f'_{b,15} = 1$  to  $\approx 43 M_{\odot}$  for  $f'_{b,15} = 0$ . Increasing  $f'_{b,15}$  decreases  $t_{seg}$ . Hence, the mass of the most massive star satisfying  $t_{seg} < t_{MS}$  increases with increasing  $f'_{b,15}$ , thus increasing  $M_{BH,max}$ . Interestingly, the higher- $f'_{b,15}$  models produce fewer BHs overall, and in particular, fewer low-mass BHs (Figure 3, top). This is because, multiple high-mass stars, each of which could potentially produce individual BHs via core-collapse, may merge to produce a single high-mass BH in high- $f'_{b,15}$  models.

Since the high-mass BHs are produced via stellar collisions at early times, we study all collisions in detail. Figure 4 shows the time and cluster-centric distance (r) of all stellar collisions (during dynamical encounters) and mergers (due to stellar evolution processes in a binary) in two representative model clusters, one with  $f'_{b,15} = 1$ (blue) and the other with  $f'_{\rm b,15}=0$  (orange). Corresponding histograms for  $t_{cl}$  and r show the distributions created using all realisations. Close to 80% (95%) of all collisions take place within the core radius,  $r_{\rm c}$  (halfmass radius,  $r_{\rm hm}$ ). This confirms the importance of quick mass segregation for high-mass stellar collisions. Each  $f'_{b,15} = 1$  model produces  $800^{+46}_{-46} (100^{+16}_{-25})$  collisions (mergers) involving at least one high-mass star. A little over half  $(449^{+23}_{-92})$  of these collisions are binary mediated. In contrast, only  $593^{+7}_{-7}$   $(4^{+1}_{-1})$  high-mass stellar collisions (mergers) occur per model for  $f'_{b,15} = 0$ . Of these collisions, only  $2^{+1}_{-1}$  per model are binary mediated.

Although, during  $t_{\rm cl}/{\rm Myr} \leq 35$ , high-mass primaries remain bound to their primordial companions most of the time (Figure 2), only  $\approx 3\%$  of the high-mass *stellar collisions* involve the primordial companions in the  $f'_{{\rm b},15} = 1$  models. In contrast, almost all ( $\approx 98\%$ ) high-mass *mergers* involve primordial companions in these models. Of course, by construct, none of the collisions/mergers in the  $f'_{{\rm b},15} = 0$  models involve primordial companions. Overall, high-mass stellar collisions/mergers seem to occur at a statistically smaller rin the models with  $f'_{{\rm b},15} = 1$  compared to the locations of those in the models with  $f'_{{\rm b},15} = 0$ . This is because




Figure 3. The total number of BHs (top), high-mass BHs (middle), and the mass of the most massive BH (bottom) formed due to core-collapse in a cluster model as a function of  $f'_{b,15}$ . Colored triangles represent individual realisations for the model clusters. Black triangles with errorbars denote the median and the quartiles for all model realisations with a given  $f'_{b,15}$ .

of the more efficient mass segregation in the higher  $f'_{\rm b,15}$  models, evident from the evolution of the 10% Lagrange radii ( $r_{\rm lagr}$ ). Although  $r_{\rm lagr}$  for all stars are comparable, the model with  $f'_{\rm b,15} = 1$  has a smaller  $r_{\rm lagr}$  for 10–100  $M_{\odot}$  stars than that for the model with  $f'_{\rm b,15} = 0$ .

Interestingly, most of the high-mass collisions/mergers occur during a short time window,  $3 \leq t_{\rm cl}/{\rm Myr} \leq 5$  independent of  $f'_{\rm b,15}$ . This is because of an enticing com-

petition between core contraction due to two-body relaxation and rapid expansion due to stellar mass loss via winds and remnant formation in young star clusters. The evolution of the central density  $(\rho_c)$  helps understand the process clearly (Figure 4). Until  $t_{\rm cl}/{\rm Myr} \lesssim 3$ ,  $\rho_c$  increases due to two-body relaxation and mass segregation. When  $t_{\rm cl}/{\rm Myr} \approx 3$ , the high-mass stars start undergoing SN. Afterwards, the mass-loss during remnant formation leads to rapid expansion of the whole cluster decreasing  $\rho_c$  by about an order of magnitude. Hence, during a small time window, high-mass stars attain their largest sizes immediately before SN and  $\rho_c$ remains sufficiently high. This leads to a sharp spike in high-mass stellar collisions. While a more efficient mass segregation leads to a higher  $\rho_c$  in the higher  $f'_{b,15}$ models, host to the same stellar populations, all models independent of  $f'_{b,15}$  exhibit the spike in high-mass stellar collisions/mergers during the same time window.

# 3.2. High-mass BHs via BH-BH Mergers

So far we have focused on the early life of a star cluster and the formation of high-mass BHs via core-collapse of collision products. However, high-mass BHs can continue to form via BH-BH mergers through dynamical processes until most BHs are ejected from the cluster (e.g., Rodriguez et al. 2018). We now shift our attention to the overall production of all BHs throughout the entire lifetime of the star cluster. In this context, when a BH merges with another BH, we consider that a new BH is formed.<sup>3</sup>

In Figure 5, we show the distributions of  $M_{\rm BH}$  for all BHs formed via any channel inside the cluster during the whole simulated lifetime,  $t_{\rm cl}/{\rm Gyr} = 13.8$ . Note that we do not consider the changes that may come to the BHs after they are ejected from the cluster, for example, via mass transfer or GW-driven mergers outside the cluster. The  $M_{\rm BH}$  distributions are very similar for all models with widely different  $f_{\rm b,15}'$ . The only statistically significant difference can be seen at low  $\log(M_{\rm BH}/M_{\odot}) \lesssim 0.5$ ; high  $f_{\rm b,15}'$  models produce fewer low-mass BHs. This is a relic of the early evolution (subsection 3.1, Figure 3).

In all models, independent of  $f'_{b,15}$ , the total number of BHs produced is dominated by core-collapse; only ~ 1–2% of all BHs form via BH-BH mergers (Figure 6, top). However, the picture changes completely when only high-mass BHs are considered. Production of high-

<sup>&</sup>lt;sup>3</sup> CMC assumes that BHs do not grow in mass during collisions with MS or giant stars (Hurley et al. 2002; Breivik et al. 2020; Rodriguez et al. 2022). BHs can grow in mass via collisions with white dwarfs, although these are incredibly rare. BHs, however, can grow via mass accretion from a binary companion.




Figure 4. All stellar collisions/mergers from example cluster models. In the main panel, we show examples from single realisations of models with  $f'_{b,15} = 1$  (blue) and 0 (orange). The colored markers and grey dots denote all collisions/mergers involving at least one high-mass star and those without any high-mass star involvement, respectively. We show the 10% Lagrange radii for all stars (dashed) and stars with mass  $10 < M_*/M_{\odot} < 100$  (dash-dot). We also show the average central density  $\rho_c$  (solid with  $1\sigma$  range denoted by the shaded regions), the scale for  $\rho_c$  is shown on the right vertical axis. The top and left panels show the corresponding distributions for high-mass stellar collision/merger times and locations from all models with  $f'_{b,15} = 1$  (blue) and 0 (orange), respectively. The hatched-shaded regions in the left panel represent the ranges of the half-mass radius ('+'-hatched) and the core radius ('x'-hatched) the cluster models attain during this early evolution.

mass BHs is dominated by BH-BH mergers for all  $f'_{b,15}$ . Moreover, the number of high-mass BHs produced via BH-BH mergers do not show any significant correlation with  $f'_{b,15}$ . Independent of  $f'_{b,15}$ , the models create  $\approx 20$ high-mass BHs via BH-BH mergers per model compared to the maximum  $\approx 4$  via core-collapse of stellar collision products (Table 1). As a result, at late times, the total number of high-mass BHs produced by a cluster overall do not show a correlation with  $f'_{b,15}$  (Figure 6, middle) despite the strong positive correlation between  $f'_{b,15}$ and high-mass BHs produced via core-collapse of stellar collision products (Figure 3). Interestingly, even the most massive BH is typically formed via BH-BH mergers for all  $f'_{b,15}$  models except  $f'_{b,15} = 1$  where  $M_{\rm BH,max}$  from core-collapse of stellar collision products win by a whisker (Figure 6, bottom; Table 1).

Figure 7 shows the mass and formation time for all new BHs in two sets of models, with  $f'_{b,15} = 1$  (blue) and 0 (orange). The dots (open circles) denote BHs formed via stellar core-collapse (BH-BH merger). The top axis shows the corresponding redshift, z assuming a typical present-day age for globular clusters,  $t_{cl}/\text{Gyr} \sim 10$  (e.g., VandenBerg et al. 2013) as a reference. As expected, high-mass BH formation via core-collapse and BH-BH mergers operate on very different timescales-  $t_{cl}/\text{Myr} \lesssim$ 10 is exclusive to stellar core-collapse whereas BH-BH mergers typically operate much later. This is because after remnant formation and the resulting expansion of




Figure 5. The same as Figure 1, except here we also consider the BHs that form via BH-BH mergers throughout the Hubble time of evolution.

the cluster, it takes time for the BHs to mass segregate to the core and partake in strong encounters to form new BBHs that can merge afterwards. Models with  $f'_{\rm b,15} =$ 0 form BHs via core-collapse almost exclusively below the upper mass-gap (only one high-mass BH formed in one of the models). In contrast, models with  $f'_{\rm b,15} = 1$ can form BHs in the upper mass gap and beyond more efficiently. At later times, high-mass BHs formed via BH-BH mergers have similar properties and formation times independent of  $f'_{\rm b,15}$ .

Another interesting trend is that the dispersion in  $M_{\rm BH}$  is significantly higher in the  $f'_{\rm b,15} = 1$  models via both channels for all times. The high dispersion stems from more stellar collisions in the  $f'_{b,15} = 1$  models which creates a more diverse population of pre-collapse highmass stars giving birth to a more diverse population of BHs. Higher  $f'_{b,15}$  also helps mix BHs with relatively more disparate masses in the core simply because it is possible to bring to the core relatively lower-mass BH progenitors as part of a binary via mass segregation. In contrast, for low  $f'_{b,15}$ , at any given time, only the most massive retained population of BHs sink to the core and take part in BH dynamics. Thus, increasing  $f'_{b \ 15}$  naturally increases the diversity of BHs a cluster can form. However, this difference in the mass distribution of BH-BH merger remnants disappear after  $t_{\rm cl}/{\rm Gyr} = 2$ , by then all differences between models with different  $f'_{\rm b,15}$ are dynamically erased (Figure 7).

Note that mass-gap BHs may also form if a BH, born with mass below but close to the mass gap, accretes sufficient mass from a companion to push it into the mass




Figure 6. The same as Figure 3, except here we also consider the BHs that form via BH-BH mergers throughout the Hubble time of evolution. Furthermore, we divide the complete population of BHs (*filled circles*) into two groups: the ones that form via core-collapse (*unfilled triangles*) and the ones that form via BH-BH mergers (*unfilled squares*).




Figure 7. The birth mass and the time of formation of all the BHs in our cluster models. Blue (orange) denotes  $f'_{b,15} = 1$  (0). We include all BHs formed in the 10 different realisations of cluster models in both the groups. The shaded region denotes the upper mass-gap. The "o" ("·") represents BHs formed via BH-BH mergers (stellar core-collapse). The top axis represents indicative redshift, z, calculated assuming cosmic time,  $t_c = 14$  Gyr represents z = 0 (Carmeli et al. 2006) and the model star clusters formed at  $t_c = 4$  Gyr, i.e., their present age is 10 Gyr.

gap. So far, we have not discussed those simply because these BHs are not high-mass BHs at the time of birth. Overall, we find that the number of high-mass BHs formed via mass transfer is  $\sim 20$  for models with  $f'_{b,15} = 1$ , comparable to the ones formed via BH-BH merger, and decreases with decreasing  $f'_{b,15}$  (Table 1). Although significant in number, especially for high  $f'_{\rm b,15}$ models, the mass for these high-mass BHs is typically just above the threshold used in our models, with median  $M_{\rm BH}/M_{\odot} \approx 46$  and  $M_{\rm BH,max}/M_{\odot} \lesssim 52$ . Necessarily in binaries, these BHs preferentially take part in BH dynamics due to larger interaction cross sections and production of merging BBHs. In subsection 3.3, we discuss how high-mass BHs influenced by mass transfer may play a role in the production of high-mass BH-BH mergers in clusters with high  $f'_{b,15}$ .

# 3.3. Demographics of Merging Binary Black Holes

Merging BBHs are of special interest because they may be detected by the GWs they emit during mergers. We now focus our attention to the BBHs that merge within a Hubble time produced in our models. For this we take into account in-cluster mergers as well as those that merge via GW radiation after being ejected from the cluster (Peters 1964). Figure 8 shows the individual pre-merger  $M_{\rm BH}$  distributions for merging BBHs. Interestingly, all models independent of  $f'_{\rm b,15}$  can produce merging BBHs in the mass gap and beyond. However, there is a clear correlation between the number of merging BBHs involving high-mass BHs and  $f'_{\rm b,15}$ . For example, the median number of high-mass merging BBHs per model increases systematically from 7 to 17 as  $f'_{\rm b,15}$ increases from 0 to 1 (Table 2).

Differences in the early production of high-mass BHs and their early mergers are primarily responsible for the positive correlation between  $f'_{\rm b,15}$  and the number of merging high-mass BBHs. Moreover, as  $f'_{\rm b,15}$  increases, so does the number of high-mass BHs formed via mass transfer (Table 1). Although there is a very significant difference in both the total number of mergers as well as the number of high-mass mergers for merger time,  $t_{\rm merge}/\text{Gyr} < 1$ , these differences reduce at later times (Figure 9). In particular, after  $t_{\rm cl}/\text{Gyr} \gtrsim 4$ , there is little statistical difference between the models with these vastly different  $f'_{\rm b,15}$ .

	Full Cluster Life										
$f_{ m b,15}'$	All		High-mass		High-mass BBH Formation Channel						
	In-cluster	Ejected	In-cluster	Ejected	Core-collapse	BBH Merger	Mass Transfer				
1	$48^{+15}_{-6}$	$72^{+12}_{-3}$	$5^{+1}_{-1}$	$10^{+2}_{-0}$	17	54	82				
0.75	$54^{+10}_{-6}$	$70^{+6}_{-3}$	$3^{+1}_{-1}$	$8^{+2}_{-2}$	15	46	52				
0.5	$49^{+7}_{-7}$	$64^{+3}_{-5}$	$4^{+1}_{-2}$	$7^{+1}_{-1}$	11	44	55				
0.25	$34^{+7}_{-1}$	$54^{+4}_{-0}$	$2^{+2}_{-1}$	$4^{+2}_{-1}$	5	51	14				
0	$30^{+3}_{-4}$	$40^{+5}_{-3}$	$2^{+1}_{-1}$	$4^{+1}_{-1}$	0	62	0				
	0 < z < 1										
1	$10^{+2}_{-2}$	$26^{+4}_{-3}$	$0^{+0}_{-0}$	$3^{+1}_{-1}$	4	16	15				
0.75	$12^{+1}_{-4}$	$26^{+4}_{-3}$	$0^{+0}_{-0}$	$2^{+2}_{-1}$	3	15	7				
0.5	$12^{+2}_{-6}$	$24^{+2}_{-2}$	$0^{+0}_{-0}$	$2^{+1}_{-1}$	2	14	7				
0.25	$8^{+3}_{-4}$	$22^{+1}_{-2}$	$0^{+0}_{-0}$	$2^{+1}_{-1}$	0	17	4				
0	$10^{+5}_{-2}$	$20^{+3}_{-3}$	$0^{+1}_{-0}$	$1^{+0}_{-0}$	0	14	0				

Table 2. The demographics of BBHs merging within a Hubble time in our models. For each  $f'_{b,15}$ , we show the numbers for in-cluster mergers and mergers after ejection from the clusters overall and those involving high-mass BHs. Furthermore, we show the contributions from different formation channels to the high-mass merging BBHs. The top part of the table shows these numbers throughout the  $\approx 14$  Gyr lifetime of the models, whereas, the bottom part shows the corresponding numbers within 0 < z < 1 (corresponds to a look-back time of  $\approx 8.4$  Gyr). All numbers with errors denote the medians and the quartiles obtained from the different realisations within models with a given  $f'_{b,15}$ . The numbers without errorbars, given for formation channels of high-mass merging BBHs are total over these realisations due to their small per-model numbers.

The picture becomes clearer if the formation channel of the high-mass BH that is participating in the merger is noticed as a function of  $t_{\text{merge}}$  (Figure 10). In the models with  $f'_{b,15} = 0$ , all high-mass BHs in the merging BBHs come from previous BH-BH mergers that remain in cluster, dynamically acquire another BH companion, and undergo higher generation mergers. In contrast, there are multiple possibilities in the models with high  $f'_{\rm b,15}$ . Here, the high-mass BHs could be created from core-collapse of stellar merger products, previous BH-BH mergers, and also accretion of mass from a non-BH companion that pushes a normal BH into the upper mass gap (Table 2). For example, in case of the  $f'_{b,15} = 1$ models, the contribution to high-mass merging BBHs from core-collapse of merger products remains below the other channels for all  $t_{\text{merge}}$ . Although the mass transfer channel can dominate the number of merging highmass BHs in the  $f'_{b,15} = 1$  models, the mass of these BHs remains close to the lower edge of the mass-gap irrespective of  $f'_{b,15}$ . We find that the median mass for all formally mass-gap BHs formed via mass transfer is  $46 M_{\odot}$ , just above the threshold used in our models, independent of  $f'_{h \ 15}$ . Given the uncertainties in the exact boundaries of the mass gap, in reality, most high-mass BHs formed via mass-transfer may not be considered as such if their mergers are observed. After  $t_{\text{merge}}/\text{Gyr} \gtrsim 6$ , the dominant formation channel for the high-mass merging BBHs is previous BH-BH mergers. Interestingly, the numbers of high-mass BHs in merging BBHs produced

via previous BH-BH mergers is insensitive to  $f'_{b,15}$  (Table 2).

Figure 11 shows the distributions for chirp mass,  $M_{\rm chirp} \equiv (M_{\rm BH1}M_{\rm BH2})^{3/5}/(M_{\rm BH1} + M_{\rm BH2})^{1/5}$  (e.g., Cutler & Flanagan 1994) and mass ratio,  $q_{\rm BH} \equiv$  $M_{\rm BH_1}/M_{\rm BH_2}$ , where  $M_{\rm BH_1} < M_{\rm BH_2}$  for merging BBHs as a function of  $t_{\text{merge}}$  for the models with  $f'_{b,15} = 0$  and 1. Clearly,  $M_{\rm chirp}$  for the merging BBHs show little difference for  $t_{\text{merge}}/\text{Gyr} \gtrsim 2$  despite the difference in  $f'_{b,15}$ . However, the models with  $f'_{b,15} = 0$  show a systematically higher  $M_{\rm chirp}$  for earlier  $t_{\rm merge}$ . When none of the high-mass stars are in binaries, such as the models with  $f'_{b,15} = 0$ , the most massive single stars sink first to the core due to mass segregation. Because of this dynamical mass filtering, the BHs produced would only find other BHs of similar high mass to interact with within the core at any given time (e.g., Breen & Heggie 2013; Morscher et al. 2013, 2015; Chatterjee et al. 2017b; Kremer et al. 2019). As a result, in the  $f'_{b,15} = 0$  models, dynamics naturally pushes  $M_{\rm chirp}$  towards higher values. In contrast, if the cluster contains significant high-mass stellar binaries, such as our models with  $f'_{b,15} = 1$ , lower-mass stars as part of binaries can also sink to the core via mass segregation at the same time with significantly highermass stars. As a result, BHs with a much broader range in  $M_{\rm BH}$  may be able to interact in the core, dynamically form BBHs, and merge. Due to the same reason, the merging BBHs tend to have a higher  $q_{\rm BH}$  in the  $f'_{b,15} = 0$  models compared to that in the models with  $f'_{b,15} = 1$  (Figure 11). Indeed, this relative difference in




Figure 8. Top: Distribution of pre-merger masses for merging BBHs from models with different  $f'_{b,15}$ . Colors, errorbars, and shaded region denote the same as in Figure 1. *Bottom*: The number of high-mass merging BHs produced as a function of  $f'_{b,15}$  over the full simulated time. Colored dots denote individual realisations. Black circles and errorbars denote the median and quartiles for all realisations for a given  $f'_{b,15}$ .

early mass segregation and the range in stellar masses that can arrive in the core between different  $f'_{\rm b,15}$  models is the key to understanding a variety of our results. For example, the same process is responsible for the higher dispersion in  $M_{\rm BH}$  for models with  $f'_{\rm b,15} = 1$  at any given formation time  $\leq 2 \,\rm Gyr$  (Figure 7). The same process is also responsible for a much wider spectrum of  $M_{\rm BH}$  in merging systems for models with  $f'_{\rm b,15} > 0$  compared to those found in the  $f'_{\rm b,15} = 0$  models (Figure 8).

### 4. SUMMARY AND CONCLUSIONS

Discovery of BBH mergers with pre-merger individual source-frame masses in or above the upper mass gap has generated wide-spread interest in thorough investi-




Figure 9. Histograms for the times of BBH mergers produced in our models, in-cluster as well as ejected, per cluster. The histograms and the shaded regions represent the mean and  $1\sigma$  between realisations. Blue (orange) represents models with  $f'_{b,15} = 1$  (0). Solid (dashed) line represents all BBH mergers (BBH mergers involving at least one high-mass BH). The top axis represents z assuming z = 0 is cosmic time  $t_c = 14$  Gyr and all model clusters formed 10 Gyr ago.

gations for the formation channels of these BHs. GKC21 showed that a young, massive, and dense star cluster with  $f_{b,15} = 1$  can produce BHs in the upper mass gap and beyond from core-collapse of high-mass stellar collision products. In contrast, they showed that similar models with  $f_{b,15} = 0$  fail to produce the high-mass BHs. We have investigated this in more detail. In addition to the difference in the initial high-mass stellar binary fraction, the GKC21 models with  $f_{b,15} = 0$  also contained  $\approx 74\%$  fewer high-mass stars compared to their models with  $f_{b,15} = 1$ . We carefully disentangle the effects of varying the fraction of high-mass stars in binaries and  $N_{15}$ . For this, we create models with identical stellar populations, but vary the fraction of high-mass stars in binaries,  $f'_{b,15} = 0$  to 1 (section 2). Moreover, we study the long-term evolution of these models focusing on the overall formation of high-mass BHs from a variety of channels. We study the time-dependent relative importance for these channels and implications for merging systems as a function of look-back time or z. We summarise our key results below.

• BH formation via stellar core-collapse completes within  $t_{\rm cl}/{\rm Myr} \lesssim 35$  in our models. Throughout  $t_{\rm cl}/{\rm Myr} < 35$ , most high-mass stars retain their primordial companions (Figure 2), and initially single high-mass stars remain single. Increasing  $f'_{\rm b,15}$  allows faster mass segregation which brings more high-mass stars to the core before they could




Figure 10. Histogram for the times of BBH mergers involving at least one high-mass BH produced per cluster. Blue (orange) represents  $f'_{b,15} = 1$  (0). Dashed, dotted, dashdotted, and solid histograms denote high-mass merging BHs produced via previous BH-BH mergers, core-collapse of stellar merger products, mass-transfer in a binary which pushes a normal BH into the upper mass gap, and all channels combined, respectively. All high-mass merging BHs form via previous BH-BH mergers in the models with  $f'_{b,15} = 0$ . In contrast, the high-mass merging BHs may be produced via all three channels in, e.g., the models with  $f'_{b,15} = 1$  (also see Table 2). Merger time is calculated from  $t_{cl} = 0$ . The top axis denotes indicative z assuming z = 0 is cosmic time  $t_c = 14$  Gyr and all model clusters formed 10 Gyr ago.

form remnants. Thus, despite containing identical  $N_{15}$ , very different populations of stars arrive to the core via mass segregation in the models with varying  $f'_{b,15}$ . This effectively alters the population of high-mass stars that can effectively collide/merge with other stars and form high-mass BHs via core collapse. As a result, the number of high-mass BHs produced as well as  $M_{\rm BH,max}$  produced via core-collapse show clear positive correlation with  $f'_{b,15}$  (Figure 1, 3).

• Most collisions/mergers take place within the cluster core (Figure 4). Although the high-mass stars spend most of their life before remnant formation with their primordial companions, most high-mass stellar collisions happen between non-primordial companions (e.g.,  $\approx 97\%$  for models with  $f'_{\rm b,15} =$  1; subsection 3.1). Collisions/mergers are also temporally concentrated between  $3 \leq t_{\rm cl}/{\rm Myr} \leq 5$  independent of  $f'_{\rm b,15}$ . This is because of the race between two-body relaxation and mass segrega-




Figure 11. Chirp mass (top) and mass ratio (bottom) as a function of  $t_{merge}$  for all BBH mergers combining all realisations from models with  $f'_{b,15} = 1$  (blue) and 0 (orange). Lines with shaded regions represent the moving average and  $1\sigma$  calculated using 101 chronologically sorted BBH mergers. The dashed parts of the lines near the two ends indicate that the moving average is calculated on a declining number of mergers.

tion, which increase  $\rho_c$ , and high-mass stellar evolution and remnant formation, which rapidly expand the cluster via mass loss and decrease  $\rho_c$ . As a result,  $\rho_c$  peaks during this time window. Furthermore, immediately before remnant formation, the stars achieve their largest physical sizes enhancing collision/merger potential.

• At later times, production of high-mass BHs is dominated by BH-BH mergers (Figure 5, Figure 7). The number and  $M_{\rm BH,max}$  for high-mass BHs produced via BH-BH mergers show no correlation with  $f'_{\rm b,15}$  (Figure 6). As a result, the overall number and  $M_{\rm BH,max}$  produced via all channels combined throughout the entire history of a cluster remains insensitive to  $f'_{\rm b.15}$ .

- Increasing  $f'_{b,15}$  increases the diversity of stars that can collide/merge during the early times of a cluster's life. This also increases the diversity of BHs that can interact with each other resulting in a wider spread in BH masses a cluster can produce (Figure 7). Pre-merger individual  $M_{\rm BH}$  for merging BBHs show a larger spread as  $f'_{b,15}$  increases due to the same reason (Figure 8).
- The overall number of high-mass BBH mergers show a clear correlation with  $f'_{b,15}$  (Figure 8). However, this is due to the differences in the number of merging BBHs with  $t_{merge}/Gyr < 1$ . The number of BBH mergers, both high-mass and overall, at  $t_{merge}/Gyr \gtrsim 2$  shows no statistical difference between models with varying  $f'_{b,15}$  (Figure 9). Assuming a typical present-day age of 10 Gyr for these clusters as a reference, identification of the effects of  $f'_{b,15}$  on the total produced number of merging BBHs would require sensitivity beyond z = 1.
- The mean  $M_{\rm chirp}$  for merging BBHs produced in  $f'_{\rm b,15} = 0$  models is significantly higher compared to those produced in  $f'_{\rm b,15} = 1$  models for  $t_{\rm merge}/{\rm Gyr} \lesssim 1$ . The models with  $f'_{\rm b,15}$  in between 0 and 1 show a systematic decrease in the mean  $M_{\rm chirp}$  with increasing  $f'_{\rm b,15}$ . However, these differences disappear for  $t_{\rm merge}/{\rm Gyr} > 2$  (Figure 11). The mean  $q_{\rm BH}$  for merging BBHs produced in the  $f'_{\rm b,15} = 0$  models tend to be higher compared to those produced in the  $f'_{\rm b,15} = 1$  models. This is simply because higher  $f'_{\rm b,15}$  facilitates production of and interactions between BHs with a wider spectrum in  $M_{\rm BH}$  (Figure 8), especially during the early evolution of a cluster.

In this study, we perform controlled experiments to clearly identify the effects of  $f'_{b,15}$  on the production of high-mass BHs in dense star clusters. We find that increasing  $f'_{b,15}$  enhances the rate of high-mass stellar collisions/mergers enabling the production of high-mass

BHs via core-collapse of merger products. However, we find that the effects are nuanced and the rich physics in star clusters produce trends that are seemingly contradictory. For example, although the production of high-mass BHs via core-collapse of high-mass collision products is strongly correlated with  $f'_{b,15}$ , the overall production of high-mass BHs is dominated by BH-BH mergers over the full lifetime of a cluster. Since BH-BH mergers show no correlations with  $f'_{\rm b,15}$ , the overall number of high-mass BHs a cluster can produce show no significant correlation with  $f'_{b,15}$ . While during early evolution, most high-mass stars remain bound to their primordial companions or remain single, only  $\approx 3\%$  of high-mass collisions happen between primordial companions. While the overall number of merging BBHs a cluster produces show a positive correlation with  $f'_{\rm h,15}$ , this effect stays limited to  $t_{\rm merge}/{\rm Gyr} \lesssim 1$ ; varying  $f'_{b,15}$  shows no effect in the number of merging BBHs for any  $t_{\rm merge}/{\rm Gyr} \gtrsim 2$ . While higher  $f'_{\rm b,15}$ models produce higher numbers of high-mass BHs and higher  $M_{\rm BH,max}$  during the early evolution of the cluster, the mean  $M_{\rm chirp}$  as well as  $q_{\rm BH}$  are lower for any  $t_{\rm merge}/{\rm Gyr} \lesssim 1$ . These results highlight the richness in the interrelated physical processes and outcomes star cluster dynamics studies can reveal. We also highlight the importance of expanding the discovery horizon to higher redshifts by the existing and planned GW detectors in order to probe the rich and interesting early evolution of massive and dense star clusters expected to have formed during an epoch of heightened star formation rate in the universe.

AK acknowledges support from TIFR's graduate fellowship. SC acknowledges support from the Department of Atomic Energy, Government of India, under project no. 12-R&D-TFR-5.02-0200 and RTI 4002. All simulations were performed on a cloud-based high-performance computing solution on Azure.

Software: CMC (Rodriguez et al. 2022), COSMIC (Breivik et al. 2020), cmctoolkit (Rui et al. 2021), Anaconda (Anaconda 2016), NumPy (Harris et al. 2020), pandas (pandas development team 2020; Wes McKinney 2010), matplotlib (Hunter 2007), Astropy (Astropy Collaboration et al. 2022).

### REFERENCES

Abbott, R., Abbott, T. D., Abraham, S., et al. 2020, Phys.

Rev. Lett., 125, 101102,

doi: 10.1103/PhysRevLett.125.101102

- Abbott, R., Abbott, T. D., Acernese, F., et al. 2023, Phys. Rev. X, 13, 041039, doi: 10.1103/PhysRevX.13.041039
- —. 2024, Phys. Rev. D, 109, 022001, doi: 10.1103/PhysRevD.109.022001

- Anaconda. 2016, Anaconda Software Distribution, Vers. 2-2.4.0, Anaconda Inc. https://anaconda.com/
- Askar, A., Szkudlarek, M., Gondek-Rosińska, D., Giersz, M., & Bulik, T. 2017, MNRAS, 464, L36, doi: 10.1093/mnrasl/slw177
- Astropy Collaboration, Price-Whelan, A. M., Lim, P. L., et al. 2022, ApJ, 935, 167, doi: 10.3847/1538-4357/ac7c74
- Ballone, A., Costa, G., Mapelli, M., et al. 2022, Monthly Notices of the Royal Astronomical Society, 519, 5191, doi: 10.1093/mnras/stac3752
- Banerjee, S. 2017, MNRAS, 467, 524, doi: 10.1093/mnras/stw3392
- --. 2018a, MNRAS, 473, 909, doi: 10.1093/mnras/stx2347
- —. 2018b, MNRAS, 481, 5123, doi: 10.1093/mnras/sty2608
- Banerjee, S., Baumgardt, H., & Kroupa, P. 2010, MNRAS, 402, 371, doi: 10.1111/j.1365-2966.2009.15880.x
- Barkat, Z., Rakavy, G., & Sack, N. 1967, Physical Review Letters, 18, 379, doi: 10.1103/PhysRevLett.18.379
- Breen, P. G., & Heggie, D. C. 2013, MNRAS, 432, 2779, doi: 10.1093/mnras/stt628
- Breivik, K., Coughlin, S., Zevin, M., et al. 2020, The Astrophysical Journal, doi: 10.3847/1538-4357/ab9d85
- Carmeli, M., Hartnett, J. G., & Oliveira, F. J. 2006, Foundations of Physics Letters, 19, 277, doi: 10.1007/s10702-006-0518-3
- Chatterjee, S., Fregeau, J. M., Umbreit, S., & Rasio, F. A. 2010, ApJ, 719, 915, doi: 10.1088/0004-637X/719/1/915
- Chatterjee, S., Rodriguez, C. L., Kalogera, V., & Rasio,
- F. A. 2017a, The Astrophysical Journal Letters, 836,
  L26, doi: 10.3847/2041-8213/aa5caa
- Chatterjee, S., Rodriguez, C. L., & Rasio, F. A. 2017b, The Astrophysical Journal, 834, 68, doi: 10.3847/1538-4357/834/1/68
- Costa, G., Ballone, A., Mapelli, M., & Bressan, A. 2022, Monthly Notices of the Royal Astronomical Society, 516, 1072, doi: 10.1093/mnras/stac2222
- Costa, G., Bressan, A., Mapelli, M., et al. 2021, Monthly Notices of the Royal Astronomical Society, 501, 4514, doi: 10.1093/mnras/staa3916
- Cutler, C., & Flanagan, E. E. 1994, Phys. Rev. D, 49, 2658, doi: 10.1103/PhysRevD.49.2658
- Di Carlo, U. N., Giacobbo, N., Mapelli, M., et al. 2019, Monthly Notices of the Royal Astronomical Society, 487, 2947, doi: 10.1093/mnras/stz1453
- Di Carlo, U. N., Mapelli, M., Giacobbo, N., et al. 2020, Monthly Notices of the Royal Astronomical Society, 498, 495, doi: 10.1093/mnras/staa2286
- Farag, E., Renzo, M., Farmer, R., Chidester, M. T., & Timmes, F. X. 2022, The Astrophysical Journal, 937, 112, doi: 10.3847/1538-4357/ac8b83

- Farmer, R., Renzo, M., de Mink, S. E., Fishbach, M., & Justham, S. 2020, The Astrophysical Journal, 902, L36, doi: 10.3847/2041-8213/abbadd
- Farmer, R., Renzo, M., de Mink, S. E., Marchant, P., & Justham, S. 2019, The Astrophysical Journal, 887, 53, doi: 10.3847/1538-4357/ab518b
- Fregeau, J. M., Cheung, P., Portegies Zwart, S. F., & Rasio, F. A. 2004, MNRAS, 352, 1, doi: 10.1111/j.1365-2966.2004.07914.x
- Fregeau, J. M., & Rasio, F. A. 2007, ApJ, 658, 1047, doi: 10.1086/511809
- Fryer, C. L., Belczynski, K., Wiktorowicz, G., et al. 2012, The Astrophysical Journal, 749, 91, doi: 10.1088/0004-637X/749/1/91
- Gerosa, D., & Kesden, M. 2016, Phys. Rev. D, 93, 124066, doi: 10.1103/PhysRevD.93.124066
- González, E., Kremer, K., Chatterjee, S., et al. 2021, The Astrophysical Journal, 908, L29, doi: 10.3847/2041-8213/abdf5b
- Goswami, S., Umbreit, S., Bierbaum, M., & Rasio, F. A. 2012, ApJ, 752, 43, doi: 10.1088/0004-637X/752/1/43
- Harris, C. R., Millman, K. J., van der Walt, S. J., et al. 2020, Nature, 585, 357, doi: 10.1038/s41586-020-2649-2
- Heger, A., Fryer, C. L., Woosley, S. E., Langer, N., & Hartmann, D. H. 2003, The Astrophysical Journal, 591, 288, doi: 10.1086/375341
- Heger, A., & Woosley, S. E. 2002, The Astrophysical Journal, 567, 532, doi: 10.1086/338487
- Hobbs, G., Lorimer, D. R., Lyne, A. G., & Kramer, M. 2005, Monthly Notices of the Royal Astronomical Society, 360, 974, doi: 10.1111/j.1365-2966.2005.09087.x
- Hunter, J. D. 2007, Computing in Science & Engineering, 9, 90, doi: 10.1109/MCSE.2007.55
- Hurley, J. R., Tout, C. A., & Pols, O. R. 2002, Monthly Notices of the Royal Astronomical Society, 329, 897, doi: 10.1046/j.1365-8711.2002.05038.x
- Ivanova, N., Heinke, C. O., Rasio, F. A., et al. 2006, Monthly Notices of the Royal Astronomical Society, 372, 1043, doi: 10.1111/j.1365-2966.2006.10876.x
- Jeans, J. H. 1919, Monthly Notices of the Royal Astronomical Society, 79, 408, doi: 10.1093/mnras/79.6.408
- Joshi, K. J., Nave, C. P., & Rasio, F. A. 2001, ApJ, 550, 691, doi: 10.1086/319771
- Joshi, K. J., Rasio, F. A., & Portegies Zwart, S. 2000, ApJ, 540, 969, doi: 10.1086/309350
- King, I. R. 1966, AJ, 71, 64, doi: 10.1086/109857

- Kremer, K., Ye, C. S., Chatterjee, S., Rodriguez, C. L., & Rasio, F. A. 2019, Proceedings of the International Astronomical Union, 14, 357–366, doi: 10.1017/S1743921319007269
- Kremer, K., Ye, C. S., Rui, N. Z., et al. 2020, ApJS, 247, 48, doi: 10.3847/1538-4365/ab7919
- Kremer, K., Spera, M., Becker, D., et al. 2020, The Astrophysical Journal, 903, 45, doi: 10.3847/1538-4357/abb945
- Kroupa, P. 2001, Monthly Notices of the Royal Astronomical Society, 322, 231, doi: 10.1046/j.1365-8711.2001.04022.x
- Lombardi, Jr., J. C., Proulx, Z. F., Dooley, K. L., et al. 2006, The Astrophysical Journal, 640, 441, doi: 10.1086/499938
- Marchant, P., Renzo, M., Farmer, R., et al. 2019, The Astrophysical Journal, 882, 36, doi: 10.3847/1538-4357/ab3426
- Martinez, M. A. S., Fragione, G., Kremer, K., et al. 2020, ApJ, 903, 67, doi: 10.3847/1538-4357/abba25
- Morscher, M., Pattabiraman, B., Rodriguez, C., Rasio, F. A., & Umbreit, S. 2015, ApJ, 800, 9, doi: 10.1088/0004-637X/800/1/9
- Morscher, M., Umbreit, S., Farr, W. M., & Rasio, F. A. 2013, ApJL, 763, L15, doi: 10.1088/2041-8205/763/1/L15
- Nitz, A. H., Kumar, S., Wang, Y.-F., et al. 2023, The Astrophysical Journal, 946, 59, doi: 10.3847/1538-4357/aca591
- Offner, S. S. R., Moe, M., Kratter, K. M., et al. 2023, in Astronomical Society of the Pacific Conference Series, Vol. 534, Protostars and Planets VII, ed. S. Inutsuka, Y. Aikawa, T. Muto, K. Tomida, & M. Tamura, 275, doi: 10.48550/arXiv.2203.10066
- pandas development team, T. 2020, pandas-dev/pandas: Pandas, latest, Zenodo, doi: 10.5281/zenodo.3509134
- Pattabiraman, B., Umbreit, S., Liao, W.-k., et al. 2013, ApJS, 204, 15, doi: 10.1088/0067-0049/204/2/15
- Peters, P. C. 1964, Phys. Rev., 136, B1224, doi: 10.1103/PhysRev.136.B1224
- Prieto, E. G., Kremer, K., Fragione, G., et al. 2022, The Astrophysical Journal, 940, 131, doi: 10.3847/1538-4357/ac9b0f
- Rakavy, G., & Shaviv, G. 1967, ApJ, 148, 803, doi: 10.1086/149204
- Renzo, M., Cantiello, M., Metzger, B. D., & Jiang, Y.-F. 2020, The Astrophysical Journal Letters, 904, L13, doi: 10.3847/2041-8213/abc6a6
- Rodriguez, C. L., Amaro-Seoane, P., Chatterjee, S., & Rasio, F. A. 2018, Phys. Rev. Lett., 120, 151101, doi: 10.1103/PhysRevLett.120.151101

- Rodriguez, C. L., Chatterjee, S., & Rasio, F. A. 2016, Phys. Rev. D, 93, 084029, doi: 10.1103/PhysRevD.93.084029
- Rodriguez, C. L., Morscher, M., Pattabiraman, B., et al. 2015, Phys. Rev. Lett., 115, 051101,
- doi: 10.1103/PhysRevLett.115.051101
- Rodriguez, C. L., Zevin, M., Amaro-Seoane, P., et al. 2019,
  Phys. Rev. D, 100, 043027,
  doi: 10.1103/PhysRevD.100.043027
- Rodriguez, C. L., Kremer, K., Grudić, M. Y., et al. 2020, ApJL, 896, L10, doi: 10.3847/2041-8213/ab961d
- Rodriguez, C. L., Weatherford, N. C., Coughlin, S. C., et al. 2022, The Astrophysical Journal Supplement Series, 258, 22, doi: 10.3847/1538-4365/ac2edf
- Rui, N. Z., Kremer, K., Weatherford, N. C., et al. 2021, NicholasRui/cmctoolkit: First release, 1.0, Zenodo, Zenodo, doi: 10.5281/zenodo.4579950
- Samsing, J., Askar, A., & Giersz, M. 2018, ApJ, 855, 124, doi: 10.3847/1538-4357/aaab52
- Sana, H., de Mink, S. E., de Koter, A., et al. 2012, Science, 337, 444, doi: 10.1126/science.1223344
- Sedda, M. A., Kamlah, A. W., Spurzem, R., et al. 2023, Monthly Notices of the Royal Astronomical Society, 526, 429, doi: 10.1093/mnras/stad2292
- Spera, M., Mapelli, M., Giacobbo, N., et al. 2019, Monthly Notices of the Royal Astronomical Society, 485, 889, doi: 10.1093/mnras/stz359
- VandenBerg, D. A., Brogaard, K., Leaman, R., & Casagrande, L. 2013, The Astrophysical Journal, 775, 134, doi: 10.1088/0004-637X/775/2/134
- Vink, J. S., Higgins, E. R., Sander, A. A. C., & Sabhahit, G. N. 2021, Monthly Notices of the Royal Astronomical Society, 504, 146, doi: 10.1093/mnras/stab842
- Weatherford, N. C., Fragione, G., Kremer, K., et al. 2021, The Astrophysical Journal Letters, 907, L25, doi: 10.3847/2041-8213/abd79c
- Wes McKinney. 2010, in Proceedings of the 9th Python in Science Conference, ed. Stéfan van der Walt & Jarrod Millman, 56 – 61, doi: 10.25080/Majora-92bf1922-00a
- Woosley, S. E., Heger, A., & Weaver, T. A. 2002, Reviews of Modern Physics, 74, 1015, doi: 10.1103/RevModPhys.74.1015
- Ye, C. S., Kremer, K., Rodriguez, C. L., et al. 2022, The Astrophysical Journal, 931, 84, doi: 10.3847/1538-4357/ac5b0b