

On Tight Components and Anti-Tight Components

Changhong Lu · Kan Wang · Xingxing Yu

Received: 4 June 2013 / Revised: 20 November 2014 / Published online: 21 February 2015
© The Author(s) 2015. This article is published with open access at Springerlink.com

Abstract A graph $G = (V, E)$ is called factor-critical if $G \neq \emptyset$ and $G - v$ has a perfect matching for every vertex $v \in V(G)$. A factor-critical graph G is tight (anti-tight, respectively) if for any $v \in V(G)$, any perfect matching M in $G - v$, and any $e \in M$, $|N(v) \cap V(e)| \neq 1$ ($|N(v) \cap V(e)| \neq 2$, respectively), where $N(v)$ denotes the neighborhood of v and $V(e)$ denotes the set of vertices incident with e . A graph G is minimally anti-tight if G is anti-tight but $G - e$ is not anti-tight for every $e \in E(G)$. In this paper, we prove that a connected graph is tight iff every block of the graph is an odd clique, and that every minimally anti-tight graph is triangle-free.

Keywords Perfect matching · Tight component · Anti-tight component

Supported in part by National Natural Science Foundation of China (Nos. 11371008 and 91230201).

C. Lu
Department of Mathematics, Shanghai Key Laboratory of PMMP,
East China Normal University, Shanghai 200241, People's Republic of China
e-mail: chlu@math.ecnu.edu.cn

K. Wang (✉)
College of Mathematics, Physics and Information Engineering, Zhejiang
Normal University, Jinhua 321004, Zhejiang, People's Republic of China
e-mail: wkan@zjnu.cn

X. Yu
School of Mathematics, Georgia Institute of Technology, Atlanta, GA 30332-0160, USA
e-mail: yu@math.gatech.edu

1 Introduction

Let G be a graph and H a component of G . For any $v \in V(G)$, we use $N_G(v)$ to denote the set of neighbors of v in G . For any $e \in E(G)$, we use $V_G(e)$ to denote the set of vertices of G incident with e . (If understood, we omit the reference to G .) If $G \neq \emptyset$ and $G - v$ has a perfect matching for every $v \in V(G)$, then G is called *factor-critical*. Factor-critical graphs have been extensively studied in the past [2,4,5,7,8]. We say that H is *odd* if $|H|$ is odd and that H is a *tight component* (respectively, *anti-tight component*) if

- H is factor-critical, and
- $|N(v) \cap V(e)| \neq 1$ (respectively, $|N(v) \cap V(e)| \neq 2$) for any $v \in V(H)$, any perfect matching M in $H - v$, and any $e \in M$.

When $H = G$, we simply say that G is tight (respectively, anti-tight).

Lee et al. [6] used tight components to solve a problem of Bollobás and Scott [1,9] about the dependence on minimum degree of bounds on judicious bisections. A *block* in a graph G is a maximal connected subgraph that contains no cut vertex. Hence, if a block of G is not 2-connected, then it must be induced by a cut edge of G . A complete subgraph of a graph is usually called a *clique*. It is easy to see that odd cliques are tight. Lee et al. [3] observed that if every block of a connected graph G is an odd clique then G is tight, and mentioned that it is not clear if every tight component is of this form. In this note, we answer this question in the affirmative.


Theorem 1 *A connected graph is tight iff every block of the graph is an odd clique.*

It is apparent that if a graph G is factor-critical and triangle-free, then G is anti-tight. But it is not true that every anti-tight graph G is triangle-free (see Fig. 1). It seems difficult to characterize the anti-tight graphs. A connected graph G is *minimally anti-tight* if G is anti-tight but $G - e$ is not anti-tight for every $e \in E(G)$. Concerning the minimally anti-tight graphs, we have

Theorem 2 *Every minimally anti-tight graph G is triangle-free.*

Let G be a graph and M a matching in G . A vertex v is called *matched* if it is incident with an edge in M , and otherwise v is called *unmatched*. An M -*alternating path* is a path in which the edges belong alternatively to $E(G) \setminus M$ and M . An M -*augmenting path* is an alternating path that starts from and ends at unmatched vertices. Let H be a subgraph of G , we use $M \Delta H$ to denote the symmetric difference of M and $E(H)$. In general, we follow the terminology in [3].

Fig. 1 An anti-tight graph with a triangle



2 Proof of Theorem 1

Lemma 1 *Let G_1 and G_2 be subgraphs of a connected graph G such that $G = G_1 \cup G_2$ and $|V(G_1) \cap V(G_2)| = 1$. Then G is tight iff both G_1 and G_2 are tight.*

Proof Let $x \in V(G_1) \cap V(G_2)$. Suppose G is tight. Let $v \in V(G_1)$. Then $G - v$ has a perfect matching, say M . Note that $|V(G_1)|$ is odd and thus the edge in M incident with x belongs to G_1 when $x \neq v$; so $M \cap E(G_1)$ is a perfect matching in $G_1 - v$. Now let M_1 be a perfect matching in $G_1 - v$. Since $G - x$ has a perfect matching, $G_2 - x$ has a perfect matching, say M_2 . Then $M := M_1 \cup M_2$ is a perfect matching in $G - v$. Thus, since G is tight, for any $e \in M_1$, $|N_G(v) \cap V(e)| \neq 1$; so $|N_{G_1}(v) \cap V(e)| = |N_G(v) \cap V(e)| \neq 1$. Hence, G_1 is tight. Similarly, we can show that G_2 is tight.

Now assume both G_1 and G_2 are tight. Let $v \in V(G)$. Then $v \in V(G_i)$ for some $i \in 1, 2$ and, without loss of generality, let $v \in V(G_1)$. Let M_1 be a perfect matching in $G_1 - v$, and M_2 a perfect matching in $G_2 - x$; so $M_1 \cup M_2$ is a perfect matching in $G - v$. Now for any perfect matching M in $G - v$, $M \cap E(G_1)$ is a perfect matching in $G_1 - v$ while $M \cap E(G_2)$ is a perfect matching in $G_2 - x$. Moreover, for any $e \in M \cap E(G_1)$, $|N_G(v) \cap V(e)| = |N_{G_1}(v) \cap V(e)| \neq 1$ (as G_1 is tight). Now suppose $e \in M \cap E(G_2)$. If $v \neq x$ then $|N_G(v) \cap V(e)| = 0 \neq 1$. So assume $v = x$. Then since G_2 is tight, $|N_G(v) \cap V(e)| = |N_{G_2}(v) \cap V(e)| \neq 1$. Hence, G is also tight. □

Lemma 2 *Let G be a 2-connected tight graph. Then G is a complete graph with odd order.*

Proof Suppose to the contrary that G is not a complete graph. Then we can find three vertices x, y, z such that $yx \in E(G)$, $yz \in E(G)$ but $xz \notin E(G)$. As $G - y$ has a perfect matching, say M , we let $xx_1 \in M$ and $zz_1 \in M$. Since $|\{x, x_1\} \cap N_G(y)| \neq 1$ and $|\{z, z_1\} \cap N_G(y)| \neq 1$, $x_1y \in E(G)$ and $z_1y \in E(G)$. Note that $G' := G - \{x, y, z\}$ has no perfect matching, otherwise, $G - \{x\}$ has a perfect matching containing yz and $|N_G(x) \cap \{y, z\}| = 1$, a contradiction. Then $x_1z_1 \notin E(G)$. If $xz_1 \in E(G)$, $G - z$ has a perfect matching $M' = (M - \{xx_1, zz_1\}) \cup \{xz_1, yx_1\}$ and $|N_G(z) \cap \{x, z_1\}| = 1$, a contradiction. Hence, $xz_1 \notin E(G)$. Similarly, $zx_1 \notin E(G)$.

Suppose $G - y$ has an M -alternating path $P' = xy_2 \cdots y_kz$. If $xy_2 \in M$ and $y_kz \in M$, then $y_2 = x_1$ and $y_k = z_1$, hence, $M \Delta yx_1y_2 \cdots y_kz$ is a perfect matching in $G - z$ and $|N_G(z) \cap \{y, x\}| = 1$, a contradiction. If $xy_2 \in M$ and $y_kz \notin M$, then $y_2 = x_1$ and $y_k \neq z_1$, clearly, $M \Delta y_2 \cdots y_kzz_1y$ is a perfect matching in $G - x$ and $|N_G(x) \cap \{z_1, y\}| = 1$, a contradiction. If $xy_2 \notin M$ and $y_kz \in M$, then we get a similar contradiction. Thus, $xy_2 \notin M$ and $y_kz \notin M$. Then $y_2 \neq x_1$ and $y_k \neq z_1$. So $M \Delta xy_2 \cdots y_kzz_1y$ is a perfect matching in $G - x_1$ and $|N_G(x_1) \cap \{z_1, y\}| = 1$, a contradiction.

Thus, $G - y$ does not contain any M -alternating path from x to z . If x_1 and z are linked by a path P_1 in $G - \{x, y\}$, let $P = xx_1 + P_1 = xx_1v_3 \cdots v_i v_{i+1} v_{i+2} \cdots z$ and assume that $xx_1v_3 \cdots v_i v_{i+1}$ is an M -alternating path, $v_{i-1}v_i \in M$ but $v_i v_{i+1}, v_{i+1} v_{i+2} \notin M$ for some i . Choose P so that $xx_1v_3 \cdots v_i v_{i+1}$ is longest and

$v_{i+1}v_{i+2} \cdots z$ is shortest. Then the matched vertex v'_{i+1} with v_{i+1} is not in P as $xx_1, v_3v_4, \dots, v_{i-1}v_i \in M$ and $v_{i+1}v_{i+2} \cdots z$ is shortest. If $v_iv'_{i+1} \notin E(G)$, then $M \Delta yx_1v_3 \cdots v_i$ is a perfect matching in $G - v_i$ and $|N_G(v_i) \cap \{v_{i+1}, v'_{i+1}\}| = 1$, a contradiction. Thus $v_iv'_{i+1} \in E(G)$. Now $xx_1v_3 \cdots v_iv'_{i+1}v_{i+1}v_{i+2} \cdots z$ is a path contradicts the choice of P , as it has a longer M -alternating section starting from x .

Hence x_1 and z are not linked by any path in $G - \{x, y\}$. Then since G is 2-connected, there is a path $P = xv_2 \cdots z$ in $G - y$ such that $x_1 \notin P$. Let $P = xv_2 \cdots v_iv_{i+1}v_{i+2} \cdots z$ such that $xv_2v_3 \cdots v_iv_{i+1}$ is an M -alternating path, $v_{i-1}v_i \in M$ but $v_iv_{i+1}, v_{i+1}v_{i+2} \notin M$. Choose P so that $xv_2 \cdots v_iv_{i+1}$ is longest and $v_{i+1}v_{i+2} \cdots z$ is shortest. Then the matched vertex v'_{i+1} with v_{i+1} is not in P as $v_{i+1} \neq x_1$ and $v_2v_3, \dots, v_{i-1}v_i \in M$ and $v_{i+1}v_{i+2} \cdots z$ is shortest. If $v_iv'_{i+1} \notin E(G)$, then $M \Delta yx_1xv_2 \cdots v_i$ is a perfect matching in $G - v_i$ and $|N_G(v_i) \cap \{v_{i+1}, v'_{i+1}\}| = 1$, a contradiction. Thus $v_iv'_{i+1} \in E(G)$. So $xv_2v_3 \cdots v_iv'_{i+1}v_{i+1}v_{i+2} \cdots z$ is a path contradicts the choice of P , as it has a longer M -alternating section starting from x . □

A simple corollary of Lemma 1 is the observation of Lee, Loh and Sudakov that if every block of a graph G is an odd clique then G is a tight component. Clearly, Lemma 1 and Lemma 2 imply Theorem 1 which says the converse.

3 Proof of Theorem 2

Assume to the contrary that G is minimally anti-tight and contains a triangle $xyzx$. If $G - xz - v$ has a perfect matching for every $v \in V(G)$, every such matching M is also a perfect matching in $G - v$. Since G is anti-tight, $|N_{G-xz}(v) \cap V(e)| \leq |N_G(v) \cap V(e)| \leq 1$ for every $e \in M$. Hence, $G - xz$ is anti-tight, a contradiction.

Thus, there is a vertex $u \in V(G)$ such that $G - xz - u$ contains no perfect matching. $G - y$ has a perfect matching M and $xz \notin M$. This implies $u \neq y$. Let \tilde{M} be a perfect matching in $G - u$. Since $G - xz - u$ contains no perfect matching, $xz \in \tilde{M}$. Let G' be the multigraph with $V(G') = V(G)$ and whose edges are edges in M and \tilde{M} . It is easy to see that $d_{G'}(u) = d_{G'}(y) = 1$ and for any $v \in V(G) - \{u, y\}$, $d_{G'}(v) = 2$. So there is a path $P = uv_1v_2 \cdots v_ly$ in G' between u and y , and the length of P is even. If $xz \notin P$, then xz is an edge of an even cycle C which is a component of G' , and $\tilde{M} \Delta C$ is a perfect matching of $G - xz - u$, a contradiction. Hence, $xz \in P$, and let $\{v_i, v_{i+1}\} = \{x, z\}$. Since $v_iv_{i+1} \notin M$, the length of $uv_1 \cdots v_i$ is odd and $C_1 := v_i \cdots v_ly + yv_i$ is an even cycle in G . Now $\tilde{M} \Delta C_1$ is a perfect matching of $G - xz - u$, a contradiction. □

Open Access This article is distributed under the terms of the Creative Commons Attribution License which permits any use, distribution, and reproduction in any medium, provided the original author(s) and the source are credited.

References

1. Bollobás, B., Scott, A.D.: Problems and results on judicious partitions. *Random Struct. Alg.* **21**, 414–430 (2002)

2. Cai, M., Favaron, O., Li, H.: $(2, k)$ -factor-critical graphs and toughness. *Graphs Combin.* **15**, 137–142 (1999)
3. Diestel, R.: *Graph Theory*. In: Graduate Texts in Mathematics. 4th ed. Springer, Berlin (2010)
4. Došlić, T.: Mycielskians and matchings. *Discuss. Math. Graph Theory* **25**, 261–266 (2005)
5. Erdős, P., Füredi, Z., Gould, R.J., Gunderson, D.S.: Extremal graphs for intersecting triangles. *J. Combin. Theory Ser. B* **64**, 89–100 (1995)
6. Lee, C., Loh, P.S., Sudakov, B.: Bisections of graphs. *J. Combin. Theory Ser. B* **103**, 599–629 (2013)
7. Liu, Y., Yan, G.: Factor-critical graphs with given number of maximum matchings. *Graphs Combin.* **24**, 563–569 (2008)
8. Lovász, L.: A note on factor-critical graphs. *Studia Sci. Math. Hungar.* **4**, 279–280 (1973)
9. Scott, A.: Judicious partitions and related problems. In: *Surveys in Combinatorics, The Proceedings of the Twentieth British Combinatorial Conference, Durham, UK*, pp. 95–117 (2005)